

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁵ : A61K 31/50, 31/535, 31/445	A2	(11) International Publication Number: WO 93/00313 (43) International Publication Date: 7 January 1993 (07.01.93)
(21) International Application Number: PCT/US92/05330 (22) International Filing Date: 26 June 1992 (26.06.92) (30) Priority data: 720,173 27 June 1991 (27.06.91) US 894,771 10 June 1992 (10.06.92) US (71) Applicant: VIRGINIA COMMONWEALTH UNIVERSITY [US/US]; MCV Station Box 568, Richmond, VA 23298-0568 (US). (72) Inventor: GLENNON, Richard, A. ; 349 Janlar Drive, Richmond, VA 23235 (US). (74) Agents: ESMOND, Robert, W. et al.; Sterne, Kessler, Goldstein & Fox, 1225 Connecticut Avenue, N.W., Suite 300, Washington, DC 20036 (US).		(81) Designated States: AU, BB, BG, BR, CA, CS, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, PL, RO, RU, SD, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: SIGMA RECEPTOR LIGANDS AND THE USE THEREOF (57) Abstract The invention relates to methods for the treatment of central nervous system disorders, gastrointestinal disorders, drug abuse, angina, migraine, hypertension and depression by administering a pharmaceutical composition comprising an effective amount of certain sigma receptor ligands to a patient in need of such treatment. The invention further relates to novel sigma receptor ligands having high binding to the sigma receptor and pharmaceutical compositions thereof. Unexpectedly, certain of the sigma receptor ligands of the present invention have selectivity for the sigma receptor over the DA, PCP and 5-HT _{1A} receptors.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

SIGMA RECEPTOR LIGANDS AND THE USE THEREOF

5

10

Field of the Invention

The invention is in the field of medicinal chemistry. In particular, the invention relates to new compounds having high binding to the sigma receptor and pharmaceutical compositions thereof. These sigma receptor ligands are useful for the treatment of central nervous system disorders.

Background of the Invention

Brain sigma receptors are the subject of intense investigation in light of the fact that sigma receptors bind many psychotropic drugs (Sonders et al., Trends Neurosci. 11:37-40 (1988)). Moreover, certain sigma receptor ligands have antipsychotic activity which suggests that sigma receptor active compounds may be used for the treatment of schizophrenia (Largent et al., Eur. J. Pharmacol. 11:345-347 (1988)).

Certain neuroleptic (i.e. antipsychotic) agents bind with very high affinity at sigma sites. Su, T., J. Pharmacol. Exp Ther. 223:284 (1982); Tam, S.W., Proc. Nat. Acad. Sci (USA) 80:6703 (1983). One agent with very high affinity for sigma sites (K_i ca 1 nM; i.e., approximately 100-fold higher affinity than N-allyl normetrazocine (NANM)) is the neuroleptic agent haloperidol. Tam, S.W. et al., Proc. Nat. Acad. Sci (USA) 81:5618 (1984). Sigma-opiates, such as NANM, bind with low affinity at typical opiate receptors but bind with significant affinity at PCP receptors.

Current neuroleptic agents are thought to produce their effects via a dopaminergic (DA) mechanism; they display very high affinities for DA binding sites. However, not all of the potent

neuroleptic agents bind at [³H]NANM-labelled sigma sites, nor do the sigma-opiates bind at DA sites. This has led to the suggestion that the sites labelled by [³H]NANM be termed sigma-sites and not sigma-opiate sites (i.e., it may simply be coincidental that the sigma opiates possess an opiate-like chemical structure). In addition, there has been speculation that agents with high affinity for sigma sites may either (a) produce psychotic effects (if they behave as agonists), or (b) produce antipsychotic effects (if they behave as antagonists). It has further been speculated that certain neuroleptic agents, such as haloperidol, produce their antipsychotic effects by both a sigma and DA mechanism. Tam, S.W. and Cook, L., Proc. Nat. Acad. Sci. (USA) 81:5618 (1984). In fact, [³H]haloperidol, in combination with spiperone (an agent with high affinity for DA sites and essentially no affinity for sigma sites) is now commonly used to label sigma sites in radioligand binding studies.

A number of researchers have studied the structure-activity relationship of sigma ligands. For example, Manallack, D.T. et al., Eur. J. Pharmacol. 144:231-235 (1987), disclose a receptor model for the phencyclidine and sigma binding sites. Manallack et al. disclose that in a recent SAR study (Largent et al., in press), sigma site affinity was shown to be enhanced by large N-alkyl substituents, e.g., benzyl or phenethyl.

Largent, B.L. et al., Mol. Pharmacol. 32:772-784 (1987), disclose a study of the structural determinants of sigma receptor affinity. In particular, Largent et al. teach that several piperidine and piperazine derivatives have sigma receptor activity. Largent et al. also disclose that affinity for the sigma receptor is markedly influenced by the N-alkyl substituents, with more lipophilic substituents affording greater affinity for the sigma receptor binding sites.

Sharkey, J. et al., Eur. J. Pharmacol. 149:171-174 (1988), studied the sigma receptor binding activity of cocaine-related compounds.

5 The literature contains a number of suggestions that the sigma receptor is not a single, homogeneous binding site. Bowen, W.D. et al., Eur. J. Pharm. 163:309-318 (1989), disclose that the effect of U.V. radiation on sigma receptor binding depended on
10 the radioligand used to assay for it. It was also demonstrated that the binding characteristics of several sigma ligands were different in membranes from certain cell lines than in guinea pig brain membranes. (Hellewell, S.B. and Bowen, W.D., Brain Res. 527:224-253 (1990); Wu, X.-Z. et al., J. Pharmacol. Exp. Ther. 257:351-359 (1991)). At least two groups have reported significantly different pharmacology for "sigma receptors" when using different radioligands to label these sites. (Itzhak, Y. et al., J. Pharmacol. Exp. Ther. 257:141-148 (1991); Karbon, E.W. et al., Eur. J. Pharm. 93:21-27 (1991)). In addition, [³H]DTG
15 binding was found to have two components in guinea pig membranes (Rothman, R.B. et al., Mol. Pharm. 39:222-232 (1991)). An overlap of sigma sites with some of the multiple sites labeled by [³H]dextromethorphan has also been described (Musacchio, J.M. et al., Life Sci. 45:1721-1732 (1989)).


20 Hellewell and Bowen, Brain Res. 527:224-253 (1990), were the first to define the characteristics of the two putative sigma receptor subtypes, named sigma-1 and sigma-2. The primary pharmacological distinction between these two sites is the affinity of the (+) isomers of the benzomorphan opiates for the
25 binding sites. These compounds, such as (+)SKF 10,047 and (+)pentazocine show nearly two orders of magnitude higher affinity for the sigma-1 site compared to the sigma-2 site. The (-) isomers of the benzomethorphans show little selectivity between these two sites. Other distinctions noted between the
30 two sites are a preponderance of the sigma-2 sites in cell lines such as NCB-20, PC12 and NG108-15 cells (Hellewell, S.B. and Bowen, W.D., Brain Res. 527:224-253 (1990); Wu, X.-Z. et al., J. Pharmacol. Exp. Ther. 257:351-359 (1991); Georg, A. and Friedl, A., J. Pharmacol. Exp. Ther. 259:479-483 (1991); Quirion, R. et al., Trends in Pharmacological Sciences 13:85-86 (1992)).
35

There has been considerable research on amphetamine and amphetamine derivatives. For example, Aldous, F.A.B., J. Med. Chem. 17:1100-1111 (1974), discloses a structure-activity study of psychotomimetic phenylalkylamines. Aldous et al. also disclose a number of halo, methyl, and methoxy substituted amphetamines.

Fuller, R.W. et al., J. Med. Chem. 14:322-325 (1971), disclose amphetamine derivatives substituted on the 3- and 4-positions of the aromatic ring with one or more chloro, fluoro, alkyl, phenoxy, alkoxy and hydroxy substituents.


Foye, W.O. et al., J. Pharm. Sci. 68:591-595 (1979), disclose heterocyclic analogues of amphetamine having 2-furyl, 2-thienyl, 3-methyl-2-phenol, 3-pyridyl, 6-methyl-2-pyridyl, 4-chlorophenyl, and 1-naphthyl rings.

Boissier, J.R. et al., Chem. Abstr. 66:46195h (1967), disclose N-benzyl amphetamine derivatives of the Formula (I):




wherein X is methyl, CF₃, methoxy, or a halogen and R is hydrogen or methyl. These compounds reportedly have anorectic activity and low toxicity. Particular compounds disclosed by Boissier et al. include N-(1-phenyl-2-propyl)-4-chlorobenzylamine, N-(1-phenyl-2-propyl)-4-methylbenzylamine, and N-(1-phenyl-2-propyl)-4-methoxybenzylamine.

Boissier, J.R. et al., Chem. Abstr. 67:21527a (1967), disclose amphetamine derivatives of the Formula (II):



wherein R^1 is hydrogen, 4-Cl, 3-Cl, or 3- CF_3 and R^2 is 2-chlorophenyl, 4-chlorophenyl, 4-bromophenyl, 3- CF_3 -phenyl, 4-tolyl, 4-methoxyphenyl, phenyl, 2-furyl, 2-tetrahydrofuryl, 2-thienyl, or 3-thienyl. Reportedly, these compounds were tested for anorexigenic activity in rats and dogs.


Osbond, J.M. *et al.*, Chem. Abstr. 69:51816c (1968), disclose N,N-bis-(omega-phenylalkyl)amines having the Formula (III):



wherein R^1 , R^2 , and R^3 are hydrogen, chloro, CF_3 , or methoxy.

Gosztanyi, T. *et al.*, J. Label. Comp. Radiopharm. 8:293-303 (1977), disclose the preparation of N-substituted omega-haloalkyl derivatives of p-chloro amphetamine. Also disclosed, is the corresponding omega-hydroxyalkyl amine.

Coutts, R.T. *et al.*, Can. J. Microbiol. 26:844-848 (1980), disclose N-substituted p-chloro amphetamines having the following Formula (IV):




wherein R^1 is 2-butanone-3-yl, 2-hydroxybutane-3-yl, 1-hydroxybutane-3-yl, or acetate.

Fuller, R.W. *et al.*, J. Pharm. Pharmacol. 25:828-829 (1973), disclose lipid-soluble derivatives of amphetamine comprising 2-chloro, 3-chloro, 4-chloro, and beta, beta-difluoro-

amphetamine, and the effect thereof on amphetamine levels in the brain.

Fuller, R.W. *et al.*, Neuropharmacology **14**:739-746 (1975), disclose 4-chloroamphetamine, 4-bromoamphetamine, and 4-fluoroamphetamine and the effect thereof on serotonin metabolism.

Conde, S. *et al.*, J. Med. Chem. **21**:978-981 (1978), disclose thiophene analogues of chloroamphetamine having the following Formula (V):




wherein X, Y, and Z are chloro or hydrogen, and the effect thereof on serotonin levels in the brain.

Lukovits, I., Int. J. Quantum. Chem. **20**:429-438 (1981), discloses various halo, methyl, and methoxy ring-substituted amphetamines, and the inhibitory potencies thereof on phenylethanolamine-N-methyl transferase.

Law, B., J. Chromatog. **407**:1-18 (1987), discloses amphetamine analogues comprising 1-methyl-2-(2'-naphthyl)ethylamine, N-isopropyl-2-(2'-naphthyl)ethylamine, and N-isopropyl-2-phenylethylamine.


Johansson, A.M. *et al.*, J. Med. Chem. **30**:602-611 (1987), disclose N-substituted 2-aminotetralins of the Formula (VI):





wherein R¹ is OH or OMe and R² and R³ are H or C₁-C₄ lower alkyl. These compounds were tested for dopamine receptor agonist and antagonist activities.

-7-

Hacksell, U. *et al.*, *J. Med. Chem.* **22**:1469-1475 (1979), disclose N-alkylated-2-aminotetralins of the Formula (VII):



10 wherein R¹ is OH or OMe, R² is lower alkyl, and R³ is lower alkyl or phenethyl. In particular, Hacksell *et al.*, disclose two aminotetralins of the Formulae (VIII) and (IX):




25 These compounds reportedly have dopamine-receptor stimulating activity.


McDermid, J.D. *et al.*, *J. Med. Chem.* **18**:362-367 (1975), disclose N-alkyl aminotetralins of the Formula (X):



wherein R¹ and R² is one of a large number or alkyl, heteroalkyl, and alkaryl groups. In particular, McDermid *et al.*, disclose two compounds of the Formula (XI) and (XII):



(XI)




(XII)

These compounds are reportedly dopamine receptor agonists.

Glennon, R.A. *et al.*, Pharmacol. Biochem. Behav. **21**:895-901 (1984), disclose that 2-aminotetralin is a conformationally restricted analog of amphetamine which is about one-half as effective as racemic amphetamine.


Beaulieu, M. *et al.*, Eur. J. Pharmacol. **105**:15-21 (1984), disclose N,N-disubstituted 2-aminotetralins of the Formula (XIII):



(XIII)

These compounds are reportedly potent D-2 dopamine receptor agonists.


Naiman, N. *et al.*, J. Med. Chem. **32**:253-256 (1989), disclose 2-(alkylamino)tetralin derivatives of the Formula (XIV):



(XIV)

wherein R is H, OMe, or OBz; R¹ is H, Me, or n-Pr; and R² is H, n-propyl, benzyl, phenethyl, or phenpropyl. These compounds reportedly bind to the 5-HT_{1A} receptor site.


Beecroft, R.A. et al., Tetrahedron 41:3853-3865 (1985), disclose N,N-disubstituted piperazines having the Formulae (XV)-(XVIII):



(XV)



(XVI)




(XVII)



where Ar = 1,2-Naphthyl or 9-Anthryl (XVIII)
and Z = H, 2-OMe or 4-OMe


Fuller, R.W. et al., J. Pharmacol. Exp. Therapeut. 218:636-641 (1981), disclose substituted piperazines having the following Formulae (XIX) and (XX):

-10-



10 which reportedly act as serotonin agonists and inhibit serotonin uptake or serotonin oxidation.

Fuller, R.W. et al., Res. Commun. Chem. Pathol. Pharmacol. 29:201-204 (1980), disclose the comparative effects on 5-hydroxyindole concentration in rat brain by p-chloroamphetamine and 1-(p-chlorophenyl)piperazine having the following Formulae (XXI) and (XXII):




25 Boissier, J. et al., Chem. Abstr. 61:10691c, disclose disubstituted piperazines having the Formula (XXIII):



35 wherein R¹ and R² are aryl and X is a straight or branched chain alkylene of C₁-C₃. The compounds are reportedly adrenolytics, hypotensors, potentiators of barbiturates, and depressants of the central nervous system.

Roessler, Chem. Abstr. 61:13328g, disclose piperazine derivatives of the Formula (XXIV):



wherein R = H or methoxy and R¹ = H, *p*-ethylphenyl, or *p*-chlorophenyl.

15 Ruschig, H *et al.*, Chem. Abstr. 53:3253e, disclose a large series of N,N-disubstituted piperazines including 1-benzyl-4-(3-chloro-4-methylphenyl)piperazine.

Shvedov, V.I. *et al.*, Chem. Abstr. 73:11806q (1970), disclose 4-(R²CH₂-CH₂ substituted)-1-phenyl-piperazines wherein R² is phenyl, 2-naphthyl, 3-indolyl, 2-methyl-3-indolyl or 2-benzimidazolyl.

20

Popov, O., Chem. Abstr. 67:54102m (1967), disclose disubstituted piperazines of the Formula (XXV):




wherein R is tolyl, *p*-methoxyphenyl, *m*-ethoxyphenyl, beta-naphthyl, *m*- or *p*-carboxylphenyl, 3,4-dimethoxyphenyl, 5-hydrindenyl, *p*-chloro-phenyl, *p*-bromophenyl, *p*-iodophenyl, 3,4-dichlorophenyl, and *m*- or *p*-nitrophenyl.


Glennon, R.A *et al.*, J. Med. Chem. 31:1968-1971 (1988), disclose various N,N-disubstituted piperazines having the Formulae (XXVI)-(XXIX):

35


-12-




(XXVI)



(XXVII)




(XXVIII)



(XXIX)

These compounds reportedly have high affinity for the 5-HT_{1A} serotonin binding site.

Prasad, R.N. *et al.*, *J. Med. Chem.* **11**:1144-1150 (1968), disclose N,N-disubstituted piperazines of the Formula (XXX):



(XXX)

wherein R¹ is phenyl or *o*-methoxyphenyl and R² is 2,4-dichlorophenyl, *o*-, *m*- or *p*-methoxyphenyl, 3,4-dimethoxyphenyl, or *m*-tolyl. These compounds are reported to be antihypertensive agents.

Despite the development of the above-mentioned derivatives, a need continues to exist for new sigma receptor ligands and for methods for the treatment of central nervous system disorders.

Summary of the Invention



The invention relates to the discovery that certain phenylalkyl-amine, aminotetralin, piperazine, piperidine and related derivatives have high binding to the sigma receptor and, unexpectedly, low binding for the PCP, DA and 5-HT_{1A} receptors. Thus, the sigma receptor ligands of the present invention can be used for the treatment of central nervous system disorders and drug abuse without the side effects of traditional neuroleptic agents which also bind to the DA and 5-HT_{1A} receptors.

The invention also relates to the discovery that certain phenylalkyl-amines, aminotetralins, piperazines, piperidines and related derivatives have selective binding for the sigma-1 binding site while others have selective binding to the sigma-2 binding site. Compounds which bind to the sigma-1 binding site are useful in treating gastrointestinal disorders, and are not associated with dystonic effects which are associated with binding to the sigma-2 binding site. In contrast, compounds which selectively bind to the sigma-2 binding site may block calcium channels. Thus, such calcium channel blocking sigma-2 receptor ligands may be used to treat psychosis, angina, migraine and hypertension.

The sigma receptor ligands of the present invention may also be employed in methods of treating or preventing depression.

In particular, the invention relates to methods of treating a human being suffering from central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, which comprises administering to said

human a therapeutically effective amount of a compound selected from the Formulae (XXXI) and (XXXII):



wherein said compound exhibits high binding activity with respect to the sigma receptor.

The invention also relates to certain novel sigma receptor ligands defined by Formulae (XXXI) and (XXXII) as well as pharmaceutical compositions comprising these novel sigma receptor ligands.

Surprisingly, the present inventor has discovered that certain N-substituted phenylalkylamines, although seemingly related to amphetamine, have activities which are, in fact, very much unlike amphetamine. Instead, the N-substituted phenylalkylamines have high affinity to the sigma receptors and low affinity to the DA and PCP receptors. In addition, certain of the sigma receptor ligands of the present invention have unexpectedly low affinity for the 5-HT_{1A} receptor. In addition, certain of the sigma receptor ligands of the present invention are highly selective for the sigma-1 binding site over the sigma-2 binding site. The discovery of such ligands having high affinity for the sigma receptors, in particular the sigma-1 receptor, and low affinity for other such receptors allows for the treatment of psychosis, drug abuse, gastrointestinal


disorders, and depression, and other conditions without untoward side effects. In contrast, compounds which are selective for the sigma-2 receptor are useful for the treatment of psychosis, hypertension, migraine and angina.

5

Description of the Preferred Embodiments

The invention relates to a method of treating a human being suffering from central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, which comprises administering to said human a therapeutically effective amount of a compound having the Formula (XXXI):

15



20

(XXXI)

wherein:

25

30

35

Ar is aryl or heteroaryl wherein aryl or heteroaryl can be substituted by hydrogen, halogen such as chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, an aryl ring fused to a substituted benzene ring, a substituted aryl ring fused to a benzene ring, a heteroaryl ring fused to a benzene ring, a substituted heteroaryl ring fused to a benzene ring, C₃-C₆ heterocycloalkyl, a C₃-C₆ heterocycloalkyl ring fused to a benzene ring, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl,

C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

R is hydrogen or C₁-C₆ alkyl;

5 R¹ is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy, amino, C₁-C₆ alkylamino or =O (a double bonded oxygen); or

R and R¹ together form a morpholino ring;

n is 0-5;

W is -(CH₂)_p- or -H H-, wherein p is 1-3;

10 X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein each r is 0-3 independently;

-(CH₂)_r-CH=CH-(CH₂)_r-;

15
$$\begin{array}{c} \text{O} \\ \parallel \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-;} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen);

20 Z is hydrogen, aryl, an aryl-substituted carboxylic acid group, heteroaryl or cycloalkyl, wherein aryl, heteroaryl and cycloalkyl can be substituted by hydrogen, halogen such as chloro, fluoro, bromo, iodo; CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted
25 aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or an ortho methylene
30 dioxy group;

wherein said compound exhibits high binding activity with respect to the sigma receptors.

35 The invention also relates to methods of treating central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, using compounds having the Formula (XXXIII):

-17-



5

wherein T is a cycloalkyl group or Ar as described above, n, R, and R¹, X and Z are defined above;


wherein said compound exhibits high binding activity with respect to the sigma receptors.

Especially preferred compounds within the scope of Formula (XXXIII) include N-phenethyl-1-phenyl-isopropylamine, N-phenylpropyl-1-phenylisopropylamine, N-(2-phenoxyethyl)-1-phenylisopropylamine, N-(3-phenyl-3-propanon-1-yl)-1-phenylisopropylamine, N-(4-phenylbutyl)-1-phenylisopropylamine, N-(3-(1-naphthyl)propyl)-1-phenylisopropylamine, N-(3-(2-naphthyl)propyl)-1-phenylisopropylamine, N-(3-phenyl-2-propyn-1-yl)-1-phenylisopropylamine, N-(3-phenylpropyl)-3-(4-hydroxyphenyl)isopropylamine, N-(3-phenylpropyl)-3-(4-methoxyphenyl)isopropylamine, N-(3-phenylpropyl)-3-(3-bromophenyl)isopropylamine, N-(3-phenylpropyl)-3-(4-bromophenyl)isopropylamine, N-(3-phenylpropyl)-3-(3,4-dichlorophenyl)isopropylamine, N-(3-phenylpropyl)-3-(4-iodophenyl)isopropylamine, N-(3-phenylpropyl)-3-(3-trifluoromethyl-phenyl)isopropylamine, N-(2-phenethyl)-N-methyl-1-phenylisopropyl-amine, N-(3-phenylpropyl)-1-phenylpropan-1-one-2-amine, N-(2-indane)-3-phenylisopropylamine, N-(2-indane)-3-phenylpropylamine, N,N-di-((3-phenyl)propyl)amine, N-(2-(1-naphthyl)ethyl)-1-phenylisopropylamine, N-(2-(2-naphthyl)ethyl)-1-phenylisopropylamine, N-(2-(1-naphthyl)propyl)-1-phenylisopropylamine, N-(2-(2-naphthyl)propyl)-1-phenylisopropylamine, N-(3-phenylpropyl)-1-phenyl-2-pentylamine, N-(3-phenylbutyl)-1-phenyl-2-butylamine, N,N-di-(2-ethyl-phenyl)methylamine, N,N-dibenzylamine, N-(3-phenylpropyl)-N-(6-phenylhexyl)amine, N-(3-phenylpropyl)-N-(5-phenylpentyl)amine, N-propyl-N-methyl-5-phenylpentylamine, N-methyl-N-(3-phenylpropyl)-1-isopropylamine, N-methyl-N-(3-methyl-2-butenyl)-1-isopropylamine, N-methyl-N-(3-methylbutyl)-1-isopropylamine, N-

35

methyl-N-(3-phenylbutyl)-1-phenyl-2-pentylamine, N-propyl-N-(3-phenyl)propyl)-1-phenyl-2-propylamine, N-benzyl-N-(3-phenyl)propyl)-1-phenyl-2-propylamine, N-phenyl-(5-phenyl)pentylamine, N-methyl-N-(3-phenyl)propyl-5-phenylpentylamine, N-(2-(o-methylphenyl)ethyl)-5-phenylpentylamine, N-(2-(m-methylphenyl)ethyl)-5-phenylpentylamine, N-(2-(p-methylphenyl)ethyl)-5-phenylpentylamine, N-benzyl-5-phenylpentylamine, N-benzyl-N-methyl-5-phenylpentylamine, N-(2-(3-hydroxyphenyl)ethyl)-5-phenylpentylamine, N-(2-(2-hydroxyphenyl)ethyl)-5-phenylpentylamine, N,N'-diethyl-2-(diphenylacetoxy)ethylamine, N,N'-diethyl-2-(fluorene-carboxy)ethylamine, N,N-Dimethyl-5-phenylpentylamine, N-Benzyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine, N-Benzyl-N-methyl-5-phenylpentylamine, N-Benzyl-5-phenylpentylamine, and N-(2-phenethyl)-N-methylpentylamine.


The invention also relates to methods of treating central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, with compounds related to Formula (XXXII) where W is $-(CH_2)_p-$ having the Formula (XXXIV):



wherein Ar, n, p, R, R¹, X and Z are as defined above; and

wherein said compound exhibits high binding activity with respect to the sigma receptors.

The invention also relates to methods of treating central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, with compounds of the Formula (XXXII):




10 wherein X and Z are as defined above and V is N or -CM-, wherein M is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, hydroxy, fluoro, chloro, bromo, trifluoromethyl, or represents one half of a double bond (with the neighboring endocyclic carbon);

15 R² is independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl; and


25 wherein said compound exhibits high binding activity with respect to the sigma receptors.

30 Preferably, the invention relates to the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, with a piperidine derivative having Formula (XXXV):



wherein R^2 , X and Z are as defined above.

The invention also relates to the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, with compounds having the Formula (XXXIX):



wherein R^2 , M, X and Z are as defined above;


15 U is selected from the group consisting of hydrogen, halogen such as chloro, fluoro, bromo, iodo; CF_3 , C_1 - C_6 alkoxy, C_2 - C_6 dialkoxymethyl, C_1 - C_6 alkyl, cyano, C_3 - C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkylthio, allyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, aralkoxy, C_2 - C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, an aryl ring fused to a substituted benzene ring, a substituted aryl ring fused to a benzene ring, a heteroaryl ring fused to a benzene ring, a substituted heteroaryl ring fused to a benzene ring, C_3 - C_6 heterocycloalkyl, a C_3 - C_6 heterocycloalkyl ring fused to a benzene ring, C_1 - C_6 alkylthio, C_1 - C_6 alkylsulfonyl, C_1 - C_6 haloalkylsulfonyl, C_1 - C_6 alkylsulfinyl, C_1 - C_6 haloalkylsulfinyl, arylthio, C_1 - C_6 haloalkoxy, amino, C_1 - C_6 alkylamino, C_2 - C_{15} dialkylamino, hydroxy, carbamoyl, C_1 - C_6 N-alkylcarbamoyl, C_2 - C_{15} N,N-dialkylcarbamoyl, nitro and C_2 - C_{15} dialkylsulfamoyl;

20
25
30 wherein said compound exhibits high binding activity with respect to the sigma receptors.

The invention also relates to a method of treating a human being suffering from central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, which comprises administering to said human a therapeutically effective amount of a compound of the Formula (XXXVa):

35

-21-




wherein:

10 R^3 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_2 - C_6 dialkoxymethyl, C_3 - C_{15} dialkylaminoalkyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, C_2 - C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl; and X, Y and Z are defined above,

15 wherein said compound exhibits a high binding activity with respect to the sigma receptors.

Especially preferred compounds within the scope of Formula (XXXVa) include N-methyl-N'-(4-phenyl-3-(E)butenyl)piperazine, N-methyl-N'-(4-phenyl-3-(Z)butenyl)-piperazine, N-methyl-N'-(4-(3-trifluoromethylphenyl)-3-(Z)butenyl)piperazine, N-methyl-N'-(4-phenylbutyl)piperazine, N-benzyl-N'-(4-phthalimidobutyl)-piperazine, N-(2-methoxyphenyl)-N'-(4-phthalimidobutyl)-piperazine, N-(5-phenylpentyl)-4-benzylpiperidine, N-(5-phenylpentyl)-4-benzyl-4-hydroxy-piperidine, N-benzyl-N'-(5-phenyl)pentylpiperazine and N,N'-di-(5-phenyl)pentylpiperazine.

25 The invention also relates to a method of treating a human being suffering from central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, which comprises administering to said human a therapeutically effective amount of a compound of the Formula (XXXVb):



40

wherein:

R^4 is hydrogen or an aryl group substituted with a group selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_2 - C_6 dialkoxymethyl, C_3 - C_{15} dialkylaminoalkyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, C_2 - C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl;

R^5 is hydrogen or hydroxy; and


X, Y and Z are defined above,

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

Examples of compounds having Formula (XXXVb) include but are not limited to N-(5-phenylpentyl)piperidine, N-(8-phenylheptyl)piperidine, N-(5-(4-methoxyphenyl)pentyl)piperidine, N-(3-phenylpropyl)piperidine, N-(5-cyclohexyl)pentylpiperidine, N-benzylpiperidine, N-(2-phenethyl)-4-hydroxy-4-phenylpiperidine, N-(2-phenethyl)-4-hydroxy-4-t-butylpiperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)piperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-phenylpiperidine, N-(5-(4-methoxyphenyl)-5-pentanon-1-yl)piperidine, N-(5-(4-methoxyphenyl)-5-pentanon-1-yl)-4-phenylpiperidine, N-(5-(4-methoxyphenyl)pentyl)-4-phenylpiperidine, N-(5-phenyl-5-pentanon-1-yl)-4-phenylpiperidine, N-(5-(4-chlorophenyl)pentyl)-4-phenylpiperidine, N-(5-(3-methoxyphenyl)-5-pentanon-1-yl)piperidine, N-(5-(3-chlorophenyl)-5-pentanon-1-yl)piperidine, N-(5-(3-chlorophenyl)-5-pentanon-1-yl)-4-phenylpiperidine, N-(5-(3-methoxyphenyl)-5-pentanon-1-yl)-4-phenylpiperidine, N-(4-(4-fluorophenyl)-4-butanon-1-yl)piperidine, N-(5-(4-fluorophenyl)-5-pentanon-1-yl)piperidine, N-(5-(4-fluorophenyl)-5-pentanon-1-yl)-4-phenylpiperidine, N-(5-(4-fluorophenyl)-5-pentanon-1-yl)-4-(3-chlorophenyl)-4-hydroxypiperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)piperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)-piperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-

4-(chlorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-5-pentanone-1-yl)-4-(chlorophenyl)piperidine
 N-(5-(3,4-dichlorophenyl)-5-pentanone-1-yl)-4-(chlorophenyl)-
 piperidine, N-(5-cyclopentylpentan-5-one-1-yl)piperidine and
 5 N-(5-(3,4-methylenedioxyphenyl)penta-2,4-dienyl)piperidine.

The invention also relates to a method of treating a human being suffering from central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, which comprises administering to said human a
 10 therapeutically effective amount of a compound of the Formula (XXXVc):




(XXXVc)

20 wherein X and Z are defined above,

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

An example of a compound having Formula XXXVc is N-(5-phenyl)pentyl-3-azabicyclo[3.2.2]nonane.

25 The invention also relates to a method of treating a human being suffering from central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, which comprises administering to said human a therapeutically effective amount of a tropane derivative of the
 30 Formula (XXXVd):




(XXXVd)

wherein R⁴, R⁵, X and Z are defined above, and

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

An example of compounds having Formula (XXXVd) include N-(5-phenyl)pentyl-4-phenyltropan-4-ol.

The invention also relates to the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, by administering a compound related to those of Formula XXXIII and having the Formula (XXXVI):




a is 1-8;

b is 1-8;

R is as defined above;

wherein said compound exhibits high binding activity with respect to the sigma receptors.

Preferably, compounds which are useful for the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, and which are within the scope of Formula (XXXI) are naphthyl derivatives having Formula (XXXVII):



wherein R, R², a and b, as defined above, may be the same or different;

wherein said compound exhibits high binding activity with respect to the sigma receptors.


Other compounds useful for the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, include morpholino derivatives having Formula (XXXVIII):



wherein Ar, X and Z are defined above;


wherein said compound exhibits high binding activity with respect to the sigma receptors.

Other compounds useful for the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, include cycloalkyl derivatives having Formula (LII)





wherein Cy is C₃-C₈ cycloalkyl and Ar, R¹, n, R, X, and Z are defined as above. Examples of compounds having Formula LII include 5-cyclohexylpentylamine, N-methyl-5-cyclohexylpentylamine, N,N-Dimethyl-5-cyclohexylpentylamine, N-cyclohexylmethyl-5-cyclohexyl-n-pentylamine, and N-cyclohexylmethyl-N-methyl-5-cyclohexyl-n-pentylamine.

Other compounds useful for the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, includes compounds of the Formula (LIII)



wherein X^1 is $-(CH_2)_r-C\equiv C-(CH_2)_r-$, wherein each r is 0-3 independently;



and R^2 , V , X , and Z are defined as above.

Other compounds useful for the treatment of central nervous system disorders, drug abuse, gastrointestinal disorders, hypertension, migraine, angina and depression, includes compounds of the Formula (LIV):




wherein R^5 and R^6 are independently a C_{1-8} alkyl group, R^7 is hydrogen or a C_{1-8} alkyl group substituted by an arylacetoxy group, and X is as defined above. Examples of such compounds include but are not limited to N,N' -diethyl-2-(diphenylacetoxy)ethylamine and N,N' -diethyl-2-(9-fluorenylcarboxy)ethylamine.

The invention is also related to the discovery that the aryl rings of compounds having Formula (LIV) may be missing, and that the compounds still retain high binding to the sigma receptor (Formula (LIV), R^7 = hydrogen). Such compounds include N,N -dimethyl- n -hexylamine and N -methyl- N -propylhexyl-amine.

The sigma receptor ligands of the present invention may exist in racemic form or in the optically active stereoisomeric form. Most preferably, the compounds exist in the S -(+) form.

The invention also relates to certain novel sigma receptor ligands and pharmaceutical compositions comprising the novel sigma receptor ligands. In particular, the invention relates to compounds having the Formula (XXXI):



(XXXI)

wherein Ar, n, R, R¹, W and Z are as defined above and

X is -(CH₂)_q-, wherein q is 3-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein each r is 0-3 independently;

-(CH₂)_r-CH=CH-(CH₂)_r-;

-(CH₂)_r-C(=O)-(CH₂)_r-;


-(CH₂)_r-Y-(CH₂)_r-, wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen);

wherein said compound exhibits high binding activity with respect to the sigma receptors.

Sigma receptor ligands having the above Formula (XXXI) wherein q is 3-6 have unexpectedly high binding to the sigma receptors (see Example 2, below). Preferably, q is 5.

The invention also relates to sigma receptor ligands having the Formula (XXXIII):



(XXXIII)

wherein T, n, R, R¹, and Z are defined above, and

X is -(CH₂)_q-, wherein q is 3-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein each r is 0-3 independently;

-(CH₂)_r-CH=CH-(CH₂)_r-;


-(CH₂)_r-C(=O)-(CH₂)_r-;

-(CH₂)_r-Y-(CH₂)_r-, wherein Y is O or S; or

C_1-C_6 alkyl (wherein Z is hydrogen);
wherein said compound exhibits high binding activity with respect
to the sigma receptors.

Sigma receptor ligands having the above Formula (XXXV)
wherein q is 3-6 also have unexpectedly high binding to the sigma
receptors (see Example 2). Most preferably, q is 5.

The invention also relates to compounds of the Formula
(XXXII):



(XXXII)

wherein R^2 , V and Z are as defined above;

X is $-(CH_2)_r-C\equiv C-(CH_2)_r-$, wherein r is 0-3;

$-(CH_2)_r-CH=CH-(CH_2)_r-$;

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$;


$-(CH_2)_r-Y-(CH_2)_r-$, wherein Y is O or S ; or

C_1-C_6 alkyl (wherein Z is hydrogen);

wherein said compound exhibits high binding activity with respect
to the sigma receptors.

Sigma receptor ligands having the above Formula (XXXII)
wherein q is 3-6 also have unexpectedly high binding to the sigma
receptors (see Example 2).

The invention also relates to compounds of the Formula
(XXXV):



(XXXV)

wherein R^2 and Z are as defined above;

X is $-(CH_2)_r-C\equiv C-(CH_2)_r-$, wherein each r is 0-3 independently;

$-(CH_2)_r-CH=CH-(CH_2)_r-$;

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$;


$-(CH_2)_r-Y-(CH_2)_r-$, wherein Y is O or S; or

C_1-C_6 alkyl (wherein Z is hydrogen);

wherein said compound exhibits high binding activity with respect to the sigma receptors.

Sigma receptor ligands having the above Formula (XXXV) also have unexpectedly high binding to the sigma receptors (see Example 2).

The invention also relates to compounds which are related to the Formula XXXII and having the Formula (XXXIX):




(XXXIX)

wherein R^2 , M, X and Z are as defined above;

U is selected from the group consisting of hydrogen, halogen such as chloro, fluoro, bromo, iodo; CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, an aryl ring fused to a substituted benzene ring, a substituted aryl ring fused to a benzene ring, a heteroaryl ring fused to a benzene ring, a substituted heteroaryl ring fused to a benzene ring, C_3-C_6 heterocycloalkyl, a C_3-C_6 heterocycloalkyl ring fused to a benzene ring, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkylamino, C_2-C_{15} dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro and C_2-C_{15} dialkylsulfamoyl;

wherein said compound exhibits high binding activity with respect to the sigma receptors.

The invention also relates to a compound of the Formula (XXXVa):



wherein:


15

R^3 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_2 - C_6 dialkoxymethyl, C_3 - C_{15} dialkylaminoalkyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, C_2 - C_6 acyl, aryl, substituted aryl, alkaryl, substituted alkaryl, aralkyl, substituted aralkyl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl; and X, Y and Z are defined above,

20

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

The invention also relates to a compound of the Formula (XXXVb):



wherein:

35

R^4 is hydrogen or an aryl group substituted with a group selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_2 - C_6 dialkoxymethyl, C_3 - C_{15} dialkylaminoalkyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, C_2 - C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl;


R^5 is hydrogen or hydroxy; and

X, Y and Z are defined above,

40

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

The invention also relates to a compound of the Formula (XXXVc):



(XXXVc)

wherein X and Y are defined above,

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

The invention also relates to a tropane derivative of the Formula (XXXVd):




(XXXVd)

wherein R^4 , R^5 , X and Z are defined above, and

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

The invention also relates to a compound of the Formula (XXXVI):



(XXXVI)

wherein

a is 1-8;


b is 1-8;

R is hydrogen or C_1 - C_6 alkyl;

R^2 is independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro and C_2-C_{15} dialkylsulfamoyl;

wherein said compound exhibits high binding activity with respect to the sigma receptors.


The invention also relates to naphthyl derivatives within the scope of Formula (XXXI) having Formula (XXXVII):



wherein R , R^1 , a and b , as defined above, may be the same or different;

wherein said compound exhibits high binding activity with respect to the sigma receptors.

The invention also relates to morpholino derivatives having Formula (XXXVIII):



wherein Ar , X and Z are defined above;


wherein said compound exhibits high binding activity with respect to the sigma receptors.

Also as derivatives of compound XXXIII, this invention is concerned with a compound having the Formula (LII)






wherein Cy is C₃-C₈ cycloalkyl and Ar, R¹, n, R, X, and Z are defined as above.

Compounds derived from the Formula XXXII are also an aspect of the present invention. These compounds include a compound of the Formula (LIII)



wherein X¹ is -(CH₂)_r-C≡C-(CH₂)_r-, wherein each r is 0-3 independently;



and R², V, X, and Z are defined as above.

The invention also relates to compounds having the Formula (LIV):



wherein R⁵ and R⁶ are independently a C₁₋₈ alkyl group, R⁷ is hydrogen or a C₁₋₈ alkyl substituted by an arylacetoxy or arylcarboxy group, and X is as defined above. Examples of compounds having Formula LIV include N,N-dimethyl-n-hexylamine, N-methyl-N-propylhexylamine, N,N'-diethyl-2-

(diphenylacetoxy)ethylamine, N,N'-diethyl-2-(fluorene-carboxy)ethylamine, N,N-diethyl-2-(diphenylacetoxy)ethylamine and N,N-diethyl-2-(9-fluorenylcarboxy)ethylamine.

5 The compounds of the invention have high binding to the sigma receptors. The sigma receptors include both the sigma-1 and sigma-2 subtypes. See Hellewell, S.B. and Bowen, W.D., Brain Res. 527:224-253 (1990); and Wu, X.-Z. et al., J. Pharmacol. Exp. Ther. 257:351-359 (1991). A sigma receptor binding assay which
10 quantitates the binding affinity of a putative ligand for both sigma sites (against ³H-DTG, which labels both sites with about equal affinity) is disclosed by Weber et al., Proc. Natl. Acad. Sci (USA) 83:8784-8788 (1986). Alternatively, [³H]pentazocine may be used to selectively label the sigma-1 binding site in a
15 binding assay. A mixture of [³H]DTG and unlabeled (+)pentazocine is used to selectively label the sigma-2 site in a binding assay. The present invention is also directed to certain ligands which are selective for the sigma-1 and sigma-2 receptors. The
20 discovery of such ligands which are selective for one of the two sigma receptor subtypes may be an important factor in identifying compounds which are efficacious in treating central nervous system disorders with minimal side effects.

 Typical C₁-C₆ alkyl groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, i-butyl, pentyl and hexyl groups.

25 Typical C₃₋₈ cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl groups.

 Typical C₂-C₆ carboxylic acyl groups include acetyl, propanoyl, i-propanoyl, butanoyl, s-butanoyl, pentanoyl and
30 hexanoyl groups.

 Typical aryl groups include phenyl, naphthyl, phenanthryl, anthracyl and fluorene groups.

 Typical aryl-substituted carboxylic acid groups include the above-mentioned carboxylic acyl groups substituted by one or more
35 aryl groups, e.g. diphenylacetoxy and fluorene-carboxy groups.

Typical alkaryl groups include the above-listed aryl groups substituted by one or more C₁-C₆ alkyl groups.

5 Typical aralkyl groups include a C₁-C₆ alkyl group substituted by one of the above-listed aryl groups, e.g. phenethyl, phenylpropyl, phenylbutyl, phenylpentyl and phenylhexyl groups as well as the branched chain isomers thereof.

Typical C₁-C₆ alkoxy carbonyl groups include carbonyl substituted by methoxy, ethoxy, propanoxy, i-propanoxy, n-butoxy, t-butoxy, i-butoxy, pentanoxy, and hexanoxy groups.

10 Typical aralkyl groups include the above-listed C₁-C₆ alkyl groups substituted by phenyl, naphthyl, phenanthryl, and anthracyl groups.

Typical C₂-C₆ alkenyl groups include vinyl, allyl, 2-butenyl, 2-pentenyl, and 2-hexenyl groups.

15 Typical C₂-C₆ alkynyl groups include acetynyl and propargyl groups.

Typical halo groups include fluorine, chlorine, bromine and iodine.

20 Typical aroyl groups include carbonyl substituted by phenyl, naphthyl, phenanthryl, and anthracyl groups.

Typical aralkanoyl groups include carbonyl substituted by the above-listed aralkyl groups.

25 Typical aralkoxy groups include the above listed C₁-C₆ alkoxy groups substituted by phenyl, naphthyl, phenanthryl, and anthracyl groups.

Typical substituted aryl groups include the above-listed aryl groups substituted by halo, hydroxy, C₁-C₆ alkoxy, amino, and the like.

30 Typical heteroaryl groups include furyl, thienyl, pyrrolyl, thiazolyl, pyridyl, pyrimidinyl, pyrizinyl, oxazolyl and phthalimido groups which may be fused to a benzene ring.

Typical substituted heteroaryl groups include the above-listed heteroaryl groups substituted by halo, C₁-C₆ alkyl and the like.

Typical C₅-C₆ heterocycloalkyl groups include tetrahydrofuranyl, tetrahydropyranyl, piperidiny, piperaziny, morpholino and pyrrolidinyl groups.

Under the binding activity studies, an IC₅₀ value of at most about 100 nM, preferably at most about 25 nM, more preferably at most 10 nM, most preferably at most 1 nM indicates a high binding affinity with respect to the sigma receptor binding sites. In the present application, the term "high affinity" is intended to mean a compound which exhibits an IC₅₀ of less than 100 nM in a sigma receptor binding assay, preferably against ³H-DTG as disclosed by Weber *et al.*, Proc. Natl. Acad. Sci (USA) 83:8784-8788 (1986), which measures the binding affinity of compounds toward both the sigma-1 and sigma-2 sites. Especially preferred sigma ligands exhibit IC₅₀ values of less than about 25 nM, more preferably less than about 10 nM, most preferably less than about 1 nM against ³H-DTG.

The inventor has unexpectedly discovered that certain of the sigma receptor ligand of the present invention exhibit enhanced selectivity to the sigma-1 binding site while other of the sigma receptor ligands exhibit enhanced selectivity to the sigma-2 binding site. Selective binding to the sigma-1 binding site is associated with various gastrointestinal effects, inhibition of contraction of the guinea pig ileum, and inhibition of acetylcholine-induced phosphoinositide response. In contrast, compounds which exhibit selective binding to the sigma-2 receptor are associated with dystonia and may block calcium channels. See Quiron *et al.*, Trends Pharm. Sci. 13:85-86 (1992); Rothman *et al.*, Mol. Pharmacol. 39:222-232 (1991). Thus, the compounds of the present invention which are selective for the sigma-1 receptor may be used for, in addition to the treatment of psychosis, the treatment or prevention of gastrointestinal disorders such as emesis, colitis and the like, without any untoward dystonia. In addition, the compounds of the present invention which are selective for the sigma-2 receptor may be used for treating psychosis and conditions which are ameliorated by calcium channel blockers, e.g. hypertension, migraine and

angina. Compounds which are selective for the sigma-2 receptor are known to produce dystonia. However, antagonists of the sigma-2 receptor are expected to be effective in treating hypertension, migraine and angina without dystonic side effects.

5 Preferably, compounds which are selective for the sigma-1 receptor compared to the sigma-2 receptor have an IC_{50} ratio of sigma-1/sigma-2 of less than about 0.1 (see Table 10). Such compounds include, but are not limited to (\pm) N-(1-phenyl-2-propyl)-4-phenylbutylamine, R(-) N-(1-phenyl-2-propyl)-3-(2-naphthyl)propylamine, (\pm) N-[1-(1'-Naphthyl)-2-propyl]-3-phenylpropylamine, 4-Hydroxy-4-phenyl-1-(3-phenylpropyl)piperidine, N-(4-phenylbutyl)phenethylamine, Di-N-[3-(2'-naphthyl)propyl]-N-methylamine, N-(4-phenylbutyl)-benzylamine, N-(5-phenylpentyl)-(4-phenyl)butylamine, N-(5-phenylpentyl)benzylamine, N-(4-phenylbutyl)-N'-benzylpiperazine, N-4(4-phenylbutyl)-N'-benzoylpiperazine, N-(3-phenylpropyl)-1-(p-ethoxyphenyl)-2-propylamine, N-(5-phenylpentyl)phenethylamine, N-(7-phenylheptyl)benzylamine, N-(7-phenylheptyl)phenethylamine, N-(5-cyclohexylpentyl)benzylamine, N-(4-phenylbutyl)-1-phenyl-2-butylamine, N-(4-phenyl-3(E)-butenyl)-N'-methylpiperazine, N-(4-phenyl-3(Z)-butenyl)-N'-methylpiperazine, N-(4-(3-trifluoromethyl)-3-(Z)-butenyl)-N-methylpiperazine, N-(4-phenyl)-N'-methylpiperazine, N-(5-phenylpentyl)-3-phenylpropylamine, N-methyl-N-propyl-5-phenylpentylamine, N-Methyl-N-(3-phenylpropyl)-1-phenylisopropylamine, N-(5-phenylpentyl)piperidine, N,N-Dimethyl-5-phenylpentylamine, 5-cyclohexylpentylamine, N-methyl-5-cyclohexylpentylamine, N,N-Dimethyl-5-cyclohexylpentylamine, and N-Benzyl-N-methyl-5-phenylpentylamine.

30 Preferably, compounds which are selective for the sigma-2 receptor compared to the sigma-1 receptor have a ratio of sigma-1/sigma-2 of greater than about 10 (see Table 10). Such compounds include, but are not limited to N-phenyl-N'-(3-(1-phthalimido)propyl)-piperazine, and N-(4-phthalimido)butyl-N'-phenylpiperazine. N-(5-phthalimido)pentyl-N'-phenylpiperazine is
35 also expected to be highly selective for sigma-2.

Surprisingly, the inventor has also discovered that the sigma receptor ligands of the present invention exhibit low affinity to the DA and PCP receptors. In addition, certain of the sigma receptor ligands of the present invention also exhibit low affinity for the 5-HT_{1A} receptor. Thus, the sigma receptor ligands of the present invention may be used for the treatment of central nervous system disorders without the untoward side effects associated with unwanted binding at the DA, PCP and/or 5-HT_{1A} receptors. By the term "low affinity" is intended a binding affinity of >100 nM, more preferably, >1000 nM in a DA, PCP or 5-HT_{1A} binding assay. Especially preferred sigma receptor ligands have high binding to the sigma receptor and low binding to the DA, PCP and/or 5-HT_{1A} receptors, as defined herein.

By the term "central nervous system disorder" is intended both psychiatric and movement dysfunctions. The selective sigma ligands of the present invention may be used to treat psychiatric disorders including psychoses, such as schizophrenia and related disorders, mania with psychotic features, major depression with psychotic features, organic psychotic disorders and other idiopathic psychotic disorders, in addition to anxiety disorders and depression. The term "schizophrenia" is intended to include any of a group of severe emotional disorders, usually of psychotic proportions, characterized by misinterpretation and retreat from reality, delusions, hallucinations, ambivalence, inappropriate affect, and withdrawn, bizarre, or regressive behavior. See Dorland's Illustrated Medical Dictionary, 26th edition, W.B. Saunders Company, Philadelphia, Pa., pp. 1171 (1981). The sigma receptor ligands of the present invention can also be used in treating movement disorders such as Parkinson's disease, tardive dyskinesia, and dystonias. See J.M. Walker et al., Pharmacol. Rev. 42:355-402 (1990), the disclosure of which is fully incorporated by reference herein.

The sigma receptor ligands of the present invention are also useful for the treatment of drug abuse. In this aspect of the invention, the compounds of the invention are administered to an individual to ameliorate symptoms of drug withdrawal or to


reduce craving for the drug, e.g. cocaine, heroin, PCP and hallucinogens.

5 As discussed above, the sigma receptor ligands of the present invention are highly selective for the sigma receptor and show low affinity for the DA and PCP receptors. Certain specific sigma receptor ligands of the present invention also bind with low affinity to 5-HT_{1A} receptors. Thus, in addition to the treatment of central nervous system disorders, the sigma selective ligands of the present invention may also be used as a pharmacological tool in an animal model for the screening of potential sigma receptor agents.

10 The sigma receptor ligands of the present invention may also be radiolabelled with, for example, ³H, ¹¹C, ¹⁴C, ¹⁸F, ¹²⁵I and ¹³¹I. In their radiolabelled form, the sigma receptor ligands of the present invention may be used for autoradiography studies of the sigma receptor sites in tissue, especially neuronal tissue.

15 The sigma receptor ligands of the present invention may be prepared by general methods of synthesis as disclosed in Example 1. For example, a sigma receptor ligand having Formula (XXXX) may be prepared by reductive amination of a compound having Formula (XXXXI) with an aldehyde having Formula (XXXXII) according to Scheme I outlined below.


Scheme I




The starting compound having Formula (XXXXI), where W = -H, may be prepared by general methods of organic synthesis. For general methods of preparing compounds of Formula (XXXXI), reference is made to Fuller, R.W. *et al.*, J. Med. Chem. 14:322-325 (1971); Foye, W.O. *et al.*, J. Pharm. Sci 68:591-595 (1979); Bossier, J.R. *et al.*, Chem. Abstr. 66:46195h and 67:21527a (1967); Aldous, F.A.B., J. Med. Chem. 17:1100-1111 (1974); Fuller, R.W. *et al.*, J. Pharm. Pharmacol. 25:828-829 (1973); Fuller, R.W. *et al.*, Neuropharmacology 14:739-746 (1975); Conde, S. *et al.*, J. Med. Chem. 21:978-981 (1978); Lukovits, I. *et al.*, Int. J. Quantum Chem. 20:429-438 (1981); and Law, B., J. Chromatog. 407:1-18 (1987), the disclosures of which are incorporated by reference herein in their entirety. The radiolabelled derivatives having Formula (XXXX) may be prepared by, for example, using a tritiated reducing agent to perform the reductive amination or by utilizing a ¹⁴C-labelled starting material.

Alternatively, where R is H, an N-substituted carboxamide of Formula (XXXXIII) may be reduced, for example, with LiAlH_4 to give the N,N-disubstituted sigma receptor ligand having the Formula (XXXX), below (see Scheme II).

Scheme II



Alternatively, where the starting compound comprises a carbonyl group, the compound having the Formula (XXXXIV) may be reduced with, for example, AlH_3 , diborane:methyl sulfide or other standard carbonyl reducing reagent to give the sigma receptor ligand having Formula (XXXX) according to Scheme III.


Scheme III

30

The sigma receptor ligands having Formula (XXXI) may be prepared by nucleophilic displacement of an electrophile (E) by the amino derivative (XXXXV) as outlined in Scheme IV. Examples of electrophiles which may be used for this purpose include halides such as Cl, Br, or I, tosylate or mesylate.

35

Scheme IV




30

Morpholino derivatives having the Formula (XXXXVI) may be prepared by reduction of a compound of the Formula (XXXXVII) with, for example, sodium borohydride to give the ring-closed morpholino derivative (XXXXVI) according to Scheme V.

35

Scheme V




45

Alternatively, where the sigma receptor ligand comprises a tetrahydropyridine ring (Formula (XXXXVIII)), the tetrahydropyridine ring may be constructed by reaction of a 2-arylpropene derivative (Formula (XXXXIX)) with an alkyl amine


(Formula (L)) and para-formaldehyde in the presence of orthophosphoric acid (Scheme VI).

Scheme VI



Reduction of the double bond of the compound having Formula (XXXXXVIII) with, for example, hydrogen and a hydrogenation catalyst such as Pd/C or Pt gives the corresponding piperidine having Formula (LI):

Scheme VII



Also included within the scope of the present invention are the optical isomers of the compounds of the invention. The optical isomers may be separated by classical resolution techniques by, for example, formation of a salt of the amino group with an optically active acid. A particularly preferred acid for this purpose is (+)-di-*p*-toluoyl-D-tartaric acid. The resulting diastereoisomeric salt may then be separated by crystallization, chromatography, or by taking advantage of the differing solubilities of the two diastereoisomeric salts. The

free base may then be isolated by treatment with a base such as aqueous ammonia and extraction with an organic solvent. Alternatively, the optical isomers may be prepared by resolution of the starting amine used to prepare the sigma ligand.

5 Also included within the scope of the present invention are the non-toxic pharmaceutically acceptable salts of the compounds of the invention. Acid addition salts are formed by mixing a solution of the sigma ligand of the invention with a solution of
10 a pharmaceutically acceptable non-toxic acid such as hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid, oxalic acid, and the like.

 In the methods of treatment of the present invention, the pharmaceutical compositions may comprise the sigma receptor
15 ligand at a unit dose level of about 0.01 to about 500 mg/kg of body weight, or an equivalent amount of the pharmaceutically acceptable salt thereof, on a regimen of 1-4 times per day. Of course, it is understood that the exact treatment level will depend upon the case history of the animal, e.g., human being,
20 that is treated. The precise treatment level can be determined by one of ordinary skill in the art without undue experimentation.

 The pharmaceutical compositions of the invention may be administered to any animal which may experience the beneficial
25 effects of the compounds of the invention. Foremost among such animals are humans, although the invention is not intended to be so limited.

 The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended
30 purpose. For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, or buccal routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the
35 recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

In addition to the sigma receptor ligands, the new pharmaceutical preparations may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Preferably, the preparations, particularly those preparations which can be administered orally and which can be used for the preferred type of administration, such as tablets, dragees, and capsules, and also preparations which can be administered rectally, such as suppositories, as well as suitable solutions for administration by injection or orally, are present at a concentration of from about 0.01 to 99 percent, together with the excipient.

The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical preparations for oral use can be obtained by combining the sigma ligand with solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

Suitable excipients are, in particular, fillers such as saccharides, for example lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, as well as binders such as starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the above-mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are

provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropymethyl-cellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

Other pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules which may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils, or liquid paraffin. In addition, stabilizers may be added.

Possible pharmaceutical preparations which can be used rectally include, for example, suppositories, which consist of a combination of one or more of the active compounds with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules which consist of a combination of the active compounds with a base. Possible base materials include, for example, liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons.

Suitable formulations for parenteral administration include aqueous solutions of the sigma ligands in water-soluble form, for example, water-soluble salts. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include

fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and adaptations of the sigma receptor ligands as well as the variety of conditions and parameters normally encountered in clinical therapy and which are obvious to those skilled in the art are within the spirit and scope of the invention.

Examples

EXAMPLE 1 PREPARATION OF SIGMA RECEPTOR LIGANDS

The sigma receptor ligands listed in Tables 1, 2 and 3 were synthesized according to one of twelve synthetic procedures (Methods A-L) disclosed immediately below.

METHODS

METHOD A: R(-)-N-(3-Phenylpropyl)-1-phenyl-2-aminopropane Hydrochloride. A mixture of hydrocinnamaldehyde (1.11 g, 8.2 mmol) and R(-)-amphetamine (0.94 g, 7.0 mmol) in MeOH (20 mL) was hydrogenated over Pt/C 5% (0.2 g) at room temperature until the theoretical amount of hydrogen was absorbed. The methanolic solution of the base was separated from the catalyst by filtration and was treated with 10% HCl until the mixture was strongly acidic. The MeOH and H₂O were removed under reduced pressure with warming to give a crude solid, which was recrystallized several times from MeOH and MEK to give 1.6 g (80%) of colorless crystals, mp 215-217°C.

METHOD B: R(-)-N-(Cyclopropylmethyl)-1-phenyl-2-aminopropane Maleate. To a suspension of LiAlH_4 (1.49 g, 38 mmol) in anhyd. ether (50 mL) was added a solution of N-(σ -methylphenylethyl)-2-cyclopropylcarboxamide (1.5 g, 74 mmol) in anhyd. Et_2O (20 mL).
5 The mixture was heated at reflux overnight, cooled to 0°C and H_2O (2.5 mL) was added in a drop wise manner. This was followed by the addition of 2N NaOH solution (2.5 mL) and then H_2O (5 mL). After the inorganic precipitate was removed by filtration, the Et_2O solution was dried (anhyd. Na_2SO_4) and treated with a
10 solution of maleic acid (1.5 g in absolute EtOH (10 mL). The product was collected by filtration, washed with Et_2O and recrystallized from 2-PrOH/ Et_2O (3x) to afford 1.3 g (59%) of fine crystals: mp $162\text{--}163^\circ\text{C}$.

METHOD C: N-(3-Phenylpropyl)-1-(4-bromophenyl)-2-aminopropane Hydrochloride. AlH_3 was prepared by the addition of AlCl_3 (0.07g, 0.5 mmol) to a suspension of LiAlH_4 (0.064 g, 1.7 mmol) in Et_2O (50 mL) at 0°C under a nitrogen atmosphere. A solution
15 of N-hydrocinnamoyl-1-(4-bromophenyl)-2-aminopropane (100 mg, 0.29 mmol) in dry Et_2O (10 mL) was added in a drop wise manner, to the AlH_3 solution at 0°C . After the addition was complete, the mixture was allowed to stir for 0.5 h at the same temperature. Excess AlH_3 was decomposed by the addition of crushed ice
20 (1 g) followed by 15% NaOH solution (2 mL). The mixture was filtered and the organic portion was separated, washed with H_2O (20, 10, 5 mL), dried (anhyd. MgSO_4) and treated with sufficient HCl gas until precipitation was complete. The precipitate was
25 collected by filtration and recrystallized from EtOH/ Et_2O (4x) to give 20 mg (19%) of a finely divided powder: mp $176\text{--}178^\circ\text{C}$.

METHOD D: N-(3-Phenylpropyl)-1-(3-trifluoromethylphenyl)-2-aminopropane Hydrochloride. A mixture of 1-(3-trifluoromethylphenyl)-2-propanone (102 mg, 0.50 mmol), 3-phenyl-
35 1-propylamine (86 mg, 0.64 mmol), glacial acetic acid (8 mg, 0.13 mmol), and MeOH (2 mL) was allowed to stir at room temperature for 0.5 h. To this mixture was added over a 4-h period sodium

borohydride (19 mg, 0.50 mmol) and the mixture was allowed to stir at room temperature for 20 h. The solvents were removed by warming under reduced pressure to give a small amount of an oil which was cooled and treated with 10% HCl. The crude product separated as a white solid, 122 mg (68%), mp 135-145°C. The crystals were dissolved in H₂O and the solution was extracted with Et₂O. The H₂O portion was separated, the H₂O was evaporated, and the crystals reformed; recrystallization from acetone gave 66 mg (37%) of colorless crystals, mp 167-169°C.

METHOD E: R(-)-N-Benzyl-1-phenyl-2-aminopropane Hydrochloride. To a mixture of R(-)-amphetamine sulfate (0.633 g, 3.4 mmol), benzaldehyde (0.547 g, 5.2 mmol), MeOH (3 mL), and glacial acetic acid (0.5 g) was added sodium cyanoborohydride (0.263 g, 4.0 mmol) at room temperature over a 1-h period. During this addition, the pH was maintained at pH 5.5-6.0 by the addition of additional glacial acetic acid (0.25 g). After stirring 20 h at room temperature, the MeOH was removed by warming under reduced pressure. The residue was treated with an excess of 10% NaOH and the product extracted into Et₂O. The product was then extracted into an excess of 10% HCl solution; the aqueous layer was decanted and the water was removed by warming under reduced pressure to give the crude product, 0.5 g (56%), mp 170-174°C. After several recrystallizations from MeOH-MEK, the colorless crystals weighed 0.46 g (51%), mp 173-175°C.

METHOD F: R(-)-N-(2-Phenoxyethyl)-1-phenyl-2-aminopropane Hydrochloride. In a 5-mL reaction vial was placed R(-)-amphetamine (0.288 g, 2.1 mmol) and 2-phenoxyethyl chloride (0.335 g, 2.1 mmol). The vial was sealed and heated at 95°C for 20 h. The reaction mixture was cooled, washed repeatedly with Et₂O to give the crude product which melted at 155-160°C. Recrystallization (3x) using MeOH and MEK gave colorless crystals that weighed 50 mg (8%), mp 178-179°C.

METHOD G: R(-)-N-(3-Phenyl-3-oxopropyl)-1-phenyl-2-aminopropane Hydrochloride. A mixture of R(-)-amphetamine hydrochloride

(0.259 g, 1.5 mmol), acetophenone (0.635 g, 5.3 mmol), paraformaldehyde (87 mg), MeOH (1.2 mL), and conc. HCl (1 drop) was placed in a 5-mL reaction vial, stirred, and heated at 65°C for 24 h. After removing the solvent using reduced pressure and warming, the reaction mixture (which became partially solid) was dissolved in H₂O (5 mL) and was extracted twice with hexane (10 mL). The aqueous portion was made basic with 10% NaOH and the product extracted into hexane (10 mL). The product was then extracted into a 10% HCl solution and the H₂O and excess HCl were removed with warming under reduced pressure to form the crude solid, mp 135-144°C. Repeated recrystallizations using MEK and acetone gave 50 mg (11% yield) of colorless crystals, mp 146-147°C.

METHOD H: (+)- α -[N-(3-phenylpropyl)amino]propiofenone Hydrochloride. A mixture of hydrocinnamaldehyde (2 g, 15 mmol) and norephedrine (2.18 g, 14.4 mmol) in EtOH (100 mL) was hydrogenated over a catalytic amount of Pd/C 10%, at room temperature. A hydrogen uptake slightly in excess of theory was obtained. The suspension was filtered and the filtrate was treated with 2N HCl (30 mL). The acid solution was evaporated to give a residual solid, which was recrystallized twice from EtOH/Et₂O to give 2.34 g (55%) of needles: mp 211-213°C.

An ice-cooled mixture of CH₂Cl₂ (20 mL) and pyridine (0.8 g, 10 mmol) was treated with dry CrO₃ (0.5 g, 5 mmol), which was added in portions within a 30-min period. The purple-brown mixture was stirred at room temperature for 2 h. A solution of the amino alcohol prepared above (0.23 g, 1.24 mmol) in CH₂Cl₂ (5 mL) was added to the pyridine-CrO₃ mixture all at once with vigorous stirring. After 15 min, the yellow organic layer was decanted from the black, sticky precipitate and extracted with 5% NaOH (100 mL). The organic portion was filtered through a bed of anhyd. Na₂SO₄ and evaporated to dryness. The oily residue was treated with conc. HCl (5 mL), warmed, and the mixture was evaporated to dryness. The solid residue was recrystallized from 2-PrOH/Et₂O (2x) to give 150 mg (40%) of the hydrochloride salt: mp 154-155°C (solidified and melted again at 170-172°C).

METHOD I: N-Benzyl-2-phenylmorpholine Hydrochloride. To a solution of N-(2-hydroxyethyl)-N-(benzoylmethyl)-benzylamine hydrochloride (0.50 g, 1.6 mmol) and MeOH (5 mL) at 5°C was added with stirring over 4 h sodium borohydride (0.213 g, 5.6 mmol).
5 The reaction was allowed to warm to room temperature over 18 h; an additional amount of sodium borohydride (70 mg, 1.8 mmol) was added, and the reaction was stirred for an additional 24 h. A small amount of H₂O (1 ml) was added to the mixture and the solvents removed by warming under reduced pressure. Water (6 mL)
10 and Et₂O (12 mL) were added to the reaction mixture; the ether layer was removed and the Et₂O evaporated under reduced pressure to give the crude liquid product as the free base (0.42 g). Treatment of the free base with cold conc. HCl (1 mL) followed by warming under reduced pressure to remove the H₂O and excess HCl
15 afforded the crude product hydrochloride. Recrystallization from acetone gave colorless crystals of the amine hydrochloride 0.38 g (77%), mp 140-143°C.

A portion of these crystals (35 mg, 0.11 mmol) was heated at 125°C with conc. HCl (0.17 g) in a sealed reaction vial for
20 1.25 h. The H₂O and excess HCl were removed by warming under reduced pressure to give 40 mg of the crude, tan product, mp 201-205°C. Two recrystallizations from acetone gave 21 mg (66%) of the substituted morpholine hydrochloride as colorless crystals, mp 211-212°C.


25 METHOD J: 1-(4-Phenylbutyl)-4-(4-chlorophenyl)-1,2,5,6-tetrahydropyridine Maleate. To a stirred mixture of glacial acetic acid (1.9 g, 31.7 mmol) and acetic anhydride (0.4 g, 4 mmol) was added slowly (85%) orthophosphoric acid (0.43 g, 4
30 mmol). After the exothermic reaction had subsided, 4-phenylaminobutane (0.58 g, 4 mmol), paraformaldehyde (0.36 g) and 4-chloro- α -methylstyrene (0.62 g, 4 mmol) were added. The reaction mixture was stirred at 115°C for 4 h then allowed to stand at room temperature for 2 days. The mixture was diluted
35 with H₂O (10 mL), extracted with hexane and the aqueous layer made basic with Na₂CO₃. The crude product was extracted into

hexane, dried over anhyd. K_2CO_3 , and the solvent removed by evaporation. The crude free base was recrystallized from H_2O -MeOH to give the purified free base (mp 87-90°C) which was dissolved in EtOH and treated with an Et_2O solution of maleic acid to give the maleate salt. Recrystallization from EtOH- Et_2O gave 0.6 g (35% yield) of fine crystals, mp 162-164°C.

METHOD K: N-(3-Phenylpropyl)-1-(4-hydroxyphenyl)-2-aminopropane Hydrobromide. A suspension of the free base of N-(3-phenylpropyl)-1-(4-methoxyphenyl)-2-aminopropane (200 mg, 0.63 mmol) in 48% HBr was heated at reflux for 6 h. The mixture was filtered while hot and the filtrate allowed to cool to room temperature to give crystals. The crystals were collected by filtration and recrystallized from EtOH- Et_2O to give 200 mg (88% yield) of the hydrobromide salt, mp 159-160°C.

METHOD L: 1-(3-Chlorophenyl)-4-(3-phenylpropyl)piperazine Hydrochloride. To a stirred solution of diborane:dimethyl sulfide complex (2M) in THF (30 mL) was added drop wise a solution of dry THF (25 mL) and 1-(3-chlorophenyl)-4-(3-phenylpropionyl)piperazine (1.28 g, 3.86 mmol). The reaction mixture was stirred at room temperature for 18 h, quenched by the addition of MeOH-HCl (25 mL), and the solvents removed under reduced pressure. Additional MeOH was added and removed under reduced pressure to give a tan solid. This was dried for 18 h under vacuum then recrystallized from MEK to give 0.9 g (67%) of colorless crystals, mp 168-169°C.

Table 1: Properties of some Series I Compounds.



Code No.	R	R'	R''	Chiral	Method ^h	Salt	mp, °C	Yield %	As ^a	Formula	Anal ^b	Mol. Wt.
AD-51	H	H	-(CH ₂) ₃ Ph	R(-)	E	HCl	173-175	51	MK	C ₁₆ H ₂₀ ClN	o	262
AA-41	H	H	-(CH ₂) ₂ Ph	R(-)	A	HCl	184-186	49	MK	C ₁₇ H ₂₂ ClN	CHN	276
AA-67	H	H	-(CH ₂) ₂ Ph	S(+)	A	HCl	184-186	22	MK	C ₁₇ H ₂₂ ClN	CHN	276
AA-46	H	H	-(CH ₂) ₃ Ph	R(-)	A	HCl	215-217	80	MK	C ₁₈ H ₂₄ ClN	CHN	290
AD-74	H	H	-(CH ₂) ₂ -OPh	R(-)	F	HCl	178-179	8	MK	C ₁₇ H ₂₂ ClNO	CHN	292
AD-67	H	H	-(CH ₂) ₂ -COPh	R(-)	D	HCl	146-147	11	A	C ₁₈ H ₂₂ ClNO	CHN	304
AM-75	H	H	-(CH ₂) ₄ Ph	(±)	A	HCl	158-159	30	EO	C ₁₉ H ₂₂ ClNO	d	304
AM-134	H	H	-(CH ₂) ₃ -(NAPH-1) ^g	R(-)	B	HCl	165-167	29	I	C ₂₂ H ₂₆ ClN O.5H ₂ O	CHN	345
AM-140	H	H	-(CH ₂) ₃ -(NAPH-2) ^f	R(-)	B	MeI ^g	188-190	45	EO	C ₂₆ H ₂₉ ClN	CHN	420
AM-74	H	H	-CH ₂ -CH ₂ -	(±)	B	MeI	162-163	59	IO	C ₁₇ H ₂₃ NO ₄	CHN	305
AM-87	H	H	-CH ₂ -CEC-Ph	R(-)	F	HCl	223-224	14	C	C ₁₇ H ₂₀ ClN	CHN	266
AM-159	4-HO	H	-(CH ₂) ₃ -Ph	(±)	K	HBr	159-160	88	EO	C ₁₈ H ₂₄ BrNO	CHN	326
AM-156	4-CH ₃ O	H	-(CH ₂) ₃ -Ph	(±)	A	HCl	192-194	67	EO	C ₁₉ H ₂₆ ClNO	CHN	320
AM-79	3-Br	H	-(CH ₂) ₃ -Ph	(±)	C	HCl	184-185	35	IO	C ₁₈ H ₂₃ BrClN	CHN	369
AM-72	4-Br	H	-(CH ₂) ₃ -Ph	(±)	C	HCl	176-178	19	EO	C ₁₈ H ₂₃ BrClN	CHN	369
JN-152	3,4-Cl ₂	H	-(CH ₂) ₃ -Ph	(±)	C	HCl	159-161	8	H	C ₁₈ H ₂₂ Cl ₂ N	CHN	359
AM-108	4-1	H	-(CH ₂) ₃ -Ph	R(-)	E	HCl	154-155	28	EO	C ₁₈ H ₂₃ Cl ₂ N	CHN	416
AA-111	3-CF ₃	H	-(CH ₂) ₃ -Ph	(±)	D	HCl	167-169	37	H	C ₁₉ H ₂₃ ClF ₃ N	CHN	358
AA-55	H	CH ₃	-(CH ₂) ₂ -Ph	S(+)	A	HCl	184-186	56	MK	C ₁₈ H ₂₄ ClN	CHN	290

gas = Recrystallization solvents: ethyl acetate (A), acetonitrile (C), absolute ethanol (E), 2-propanol (I), 2-butanone (K), methanol (M), acetone (N), diethyl ether (O).

bANAL = Microanalysis. All new compounds were submitted for elemental analysis and found values are within 0.4% of theory.

cBenzoate salt (mp 101-103°C) previously reported: J. Pharm. Sci. 65:1462 (1975). Compound AD-51 was converted to the benzoate salt for purpose of identification: mp 100-101°C.

dLiterature (J. Pharm. Sci. 65:1462 (1975)) mp 133°C for p-toluenesulfonate salt. A small sample of AM-75 was converted to this salt for purpose of comparison: mp 132-133°C.

eNAPH-1 = 1-Naphthyl.

fNAPH-2 = 2-Naphthyl.

gMal = maleate salt.

^hThere are other ways in which the Series I compounds might have been prepared than the method shown.

Table 2) Properties of Additional Series 1 Compounds.

Code No.	R'	n	Chiral	Method ^d	Salt	mp. ^c	Yield %	RS ^a	Formula	Anal ^b	Mol. Wt.
AM-80	H	3	(±)	H	HCl	154-155	40	10	C ₁₈ H ₂₂ ClNO	CHN	304
AM-133	H	3	(±)	A	HCl	246-247	62	EO	C ₁₈ H ₂₂ ClN	CHN	288
AM-113	H	3		A	HCl	202-203	50	E	C ₁₈ H ₂₄ ClN	c	290
AM-114	H	3	(±)	A	HCl	212-213	50	EO	C ₂₂ H ₂₆ ClN	CHN	340
AM-120	H	3	(±)	A	HCl	188-189	50	EO	C ₂₂ H ₂₆ ClN	CHN	340
MT-10	H	3	(±)	B	HCl	250-251	81 ^c	I	C ₂₀ H ₂₅ ClN	CHN	315

^a See footnote a, Table 1.^b See footnote b, Table 1.^c Literature (Chem. Abstr. 52:7812 (1955)), mp 201-202°C.^d There are other ways in which the Series 1 compounds might have been prepared than the method shown.

-57-

Table 3: Properties of Series II Compounds.

Code No.	R	D	Chiral	Method ^c	Salt	mp, °C	Yield %	RS ^a	Formula	Anal ^b	Mol. Wt.
JLH-166		2		L	HCl	195-200 (DEC)	50	K	C ₁₉ H ₂₂ ClF ₃ N ₂ · 1/2 H ₂ O	e	380
JLH-170		2		L	HCl	197-199	40	K	C ₁₈ H ₂₂ Cl ₂ N ₂ · H ₂ O	CHN	355
JLH-171		3		L	HCl	168-169	67	K I	C ₁₉ H ₂₄ Cl ₂ N ₂ · H ₂ O	CHN	369
AM-128		3		A	HCl	217-218	70	EO	C ₂₀ H ₂₆ ClN	d	316
AM-127		3	(±)	A	HCl	207-209	90	EO	C ₂₀ H ₂₆ ClNO	CHN	332
AM-129		4		J	HCl	162-164	35	EO	C ₂₃ H ₂₈ ClNO ₄	CHN	442
AD-34		1	(±)	I	HCl	211-212	49	N	C ₁₇ H ₂₀ ClNO	CHN	290

^aSee footnote a, Table 1.^bSee footnote b, Table 1.^cLiterature (Chem. Abstr. 60:12028), mp 205-207°C.^dLiterature (J. Med. Chem. 9:555 (1966)), mp 214-216°C.^eThere are other ways in which the Series II compounds might have been prepared than the method shown.

EXAMPLE 2 SIGMA, PCP AND DOPAMINE RECEPTOR BINDING ASSAYSMethods

Sigma receptor binding assays using guinea pig brain membrane homogenates and the radioligand [^3H]DTG were conducted as described by Weber et al., P.N.A.S. (USA) 83:8784-8788 (1986) which indicates binding to both the sigma-1 and sigma-2 sites. Briefly, frozen whole guinea-pig brains (Biotrol, Indianapolis, IN) were homogenized in 10 volumes (w/v) of ice-cold 320 mM sucrose using a Brinkman polytron. The homogenate was centrifuged at 1,000 x g for 20 minutes at 4°C. The supernatant was centrifuged at 20,000 x g for 20 minutes at 4°C. The resulting pellet was resuspended in 10 initial volumes of 50 mM Tris/HCl buffer at pH 7.4 and centrifuged at 20,000 x g for 20 minutes at 4°C. The resulting pellet was resuspended in 5 initial volumes ice-cold 50mM Tris/Hcl (pH .4), and the final volume was adjusted to yield a protein concentration of 3 mg/ml. Aliquots of 20-ml were stored at -70°C until used, with no detectable loss of binding.

For [^3H]DTG binding assays, the frozen membrane suspensions were thawed and diluted 1:3 in 50 mM Tris/HCl (pH 7.4). To 12 x 75 mm polystyrene test tubes were added 0.8 ml of diluted membrane suspension, 0.1 ml of [^3H]DTG (Dupont/NEN) to yield a final concentration of 1.4 nM, and 0.1 ml of unlabelled drugs or buffer. The protein concentration in the 1-ml final incubation volume was 800 ug/ml, corresponding to 32 mg of brain tissue (original wet weight) and to a tissue concentration within the linear range for specific binding. Non-specific binding was defined as that remaining in the presence of 10 uM haloperidol. Incubations were terminated after 90 minutes at room temperature by addition of 4 ml of ice-cold 50mM Tris/HCl (pH 7.4) and rapid filtration of the membrane suspension through Whatman GF/B glass-fiber filters under vacuum, using a 48-well cell harvester (Brandel). The filters were washed 2 times with 4 ml of 50 mM Tris/HCl (pH 7.4). Each filter was suspended in 10 ml Cytoscint (ICI), and radioactivity was measured by liquid scintillation

spectrometry at a counting efficiency of approximately 50%. IC₅₀ values were determined by non-linear regression analysis.

PCP receptor binding assays against ³H-MK-801 were conducted as described by Keana *et al.*, Proc. Natl. Acad. Sci (USA) **86**:5631-5635 (1989); Keana *et al.*, Life Sciences **43**:965-973 (1988). For (+) ³H-MK-801 binding, 1 nM of radioligand was incubated with about 100 ug of thawed rat brain membrane protein for 4 hr at room temperature. The assays were carried out in 5 mM Tris/acetate and were stopped by rapid filtration through Whatman GF/B or Schleicher & Schuell no. 32 glass fiber filters (presoaked in 0.05 % polyethylenamine).






Dopamine D1 and D2 receptor binding assays were performed as described by Billard *et al.*, Life Sci. **35**:1885-1893 (1984) using [³H]SCH-23390 for the D1 receptors and [³H]domperidone for the D2 receptors (Baudry *et al.*, Arch. Pharmacol. **308**:231-237 (1979)).






Rat striatal membranes were prepared from frozen tissue by Poltron homogenization in 25 volumes of ice cold Tris-EDTA buffer (50 mM Tris-HCl, 1 mM EDTA, pH 7.4 at 4°C). The homogenate was centrifuged at 48,000 x g for 10 min. at 4°C, and the pellet was resuspended in 25 volumes of the same buffer. This suspension was then incubated at 37°C for 15 min., followed by recentrifugation as before. The resulting pellet was resuspended in 267 volumes of assay buffer (50 mM Tris-HCl, 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂, pH 7.4 at 37 °C).

The binding assays were conducted with 100 uL of [³H]SCH-23390 or [³H]domperidone (to give about 1 nM final), 100 uL of buffer or drug solution, and 800 uL of membrane suspension (to give about 250 ug of striatal membrane protein per assay). The tubes were incubated at 37°C for 60 min. The assays were stopped by rapid filtration over Schleicher & Schuell #32 or Whatman GF/B glass fiber filters (presoaked in 0.5% polyethyleneimine for [³H]domperidone binding), followed by two 3 mL washes of ice cold buffer using a Brandel cell harvester. After vigorous shaking, the filter disks were counted at 52% efficiency in 5 mL of Cytoscint (ICN).






The results of these binding assays appear in Table 4.






TABLE 4

#	COMPOUND	ISO-MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	D1 IC50 (uM) (n)	D2 IC50 (uM) (n)	PCP IC50 (M)
1		R-	2.87×10^{-8}	3.80×10^{-9}	5	>10 (2)	4.65 $\pm .43$ (3)	$>10^{-5}$
2		+/-	1.25×10^{-8}	1.00×10^{-10}	4	>10 (2)	1.79 $\pm .02$ (2)	$>10^{-5}$
3		R-	8.65×10^{-6}	1.25×10^{-6}	2			$>10^{-5}$
4		R-	6.85×10^{-7}	2.50×10^{-8}	2			$>10^{-5}$
5		R-	1.705×10^{-6}	1.65×10^{-8}	2			$>10^{-5}$

#	COMPOUND	ISO- MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	PCP IC50 (M)
6		R-	1.22×10^{-7}	1.65×10^{-8}	2	$>10^{-5}$
7		R-	6.24×10^{-8}	9.00×10^{-9}	2	9.00×10^{-6}
8		R-	4.71×10^{-8}	5.75×10^{-9}	2	$>10^{-5}$
9		R-	5.84×10^{-8}	7.45×10^{-9}	2	$>10^{-5}$
10		+/-	9.69×10^{-9}	5.10×10^{-10}	2	$>10^{-5}$

-63-






#	COMPOUND	ISO-MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	D1 IC50 (uM)	D2 IC50 (uM) (n)	PCP IC50 (M)
11		R-	9.15×10^{-9}	2.75×10^{-9}	2			$>10^{-5}$
12		R-	2.35×10^{-9}	6.53×10^{-10}	4			$>10^{-5}$
13		+/-	1.22×10^{-7}	2.50×10^{-9}	2			$>10^{-5}$
14		+/-	2.85×10^{-8}	5.20×10^{-9}	2			$>10^{-5}$
15			1.445×10^{-8}	1.21×10^{-9}	4	>10 (2)	1.68 ± 3.6 (2)	$>10^{-5}$






#	COMPOUND	ISO-MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	D1 IC50 (uM)	D2 IC50 (uM) (n)	PCP IC50 (M)
16		+/-	1.83×10^{-8}	2.65×10^{-9}	2			$>10^{-5}$
17		+/-	6.60×10^{-9}	1.79×10^{-9}	2			$>10^{-5}$
18		+/-	1.08×10^{-8}	0.00	2			$>10^{-5}$
19		+/-	1.493×10^{-8}	2.95×10^{-9}	4	>10 (2)	9.32 ± 4.5 (3)	$>10^{-5}$
20		+/-	5.17×10^{-9}	1.02×10^{-9}	n=3.			$>10^{-5}$

#	COMPOUND	ISO-MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	D1 IC50 (uM) (n)	D2 IC50 (uM) (n)	PCP IC50 (M)
21		+/-	9.11×10^{-9}	6.15×10^{-10}	2			$>10^{-5}$
22		R-	1.597×10^{-8}	4.64×10^{-9}	4	>10 (2)	3.81 ± 9.7 (3)	$>10^{-5}$
23		+/-	4.10×10^{-9}	2.05×10^{-10}	2			$>10^{-5}$
24		+/-	6.91×10^{-7}	1.39×10^{-7}	2			$>10^{-5}$
25			1.01×10^{-7}	4.43×10^{-8}	2			$>10^{-5}$






No.	COMPOUND	ISOMER	SIGMA IC ₅₀ (M)	SIGMA S.E.M.	n	PCP IC ₅₀ (M)
26		S,S-	2.88×10^{-7}	2.00×10^{-9}	2	$>10^{-5}$
27		S+	3.26×10^{-8}	2.05×10^{-9}	2	$>10^{-5}$
28		R-	4.80×10^{-5}	0.00	2	$>10^{-5}$
29		R-	4.95×10^{-6}	5.00×10^{-8}	2	$>10^{-5}$
30		R-	2.10×10^{-5}	2.00×10^{-6}	2	$>10^{-5}$






-67-

#	COMPOUND	ISO- MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	PCP IC50 (M)
31		+/-	2.08×10^{-8}	6.50×10^{-10}	2	$>10^{-5}$
32			1.09×10^{-8}	2.67×10^{-9}	2	$>10^{-5}$
33			1.95×10^{-8}	5.60×10^{-9}	2	$>10^{-5}$
34		+/-	2.55×10^{-6}	4.00×10^{-8}	2	$>10^{-5}$
35		+/-	2.59×10^{-7}	1.90×10^{-8}	2	$>10^{-5}$


#	COMPOUND	ISO- MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	D1 IC50 (uM) (n)	D2 IC50 (uM) (n)	PCP IC50 (M)
36			6.85×10^{-9}	1.50×10^{-10}	2			5.00×10^{-6}
37			7.75×10^{-9}	1.45×10^{-9}	2			$>10^{-5}$
38			4.30×10^{-9}	2.10×10^{-9}	2			$>10^{-5}$
39			1.20×10^{-9}	2.00×10^{-10}	2			$>10^{-5}$
40			3.06×10^{-9} 3.18×10^{-9}	3.4×10^{-10} 3.0×10^{-11}	5 ¹ 2 ¹	>10 (2)	1.25 ± 0.06 (3)	$>10^{-5}$

¹The second set of numbers represent a second lot.

I	COMPOUND	ISO- MER	SIGMA IC ₅₀ (M)	SIGMA S.E.M.	n	D1 IC ₅₀ (uM) (n)	D2 IC ₅₀ (uM) (n)	PCP IC ₅₀ (M)
41			4.58x10 ⁻⁹ 5.59x10 ⁻⁹	8.7x10 ⁻¹⁰ 2.06x10 ⁻⁹	5 2 1	>10 (2)	1.92 ±.19 (3)	>10 ⁻⁵
42			5.20x10 ⁻⁹	2.00x10 ⁻¹⁰	2			>10 ⁻⁵
43			2.27x10 ⁻⁸	2.10x10 ⁻⁹	2			>10 ⁻⁵
44			6.90x10 ⁻⁹	4.10x10 ⁻⁹	2			>10 ⁻⁵
45			5.95x10 ⁻⁹	3.95x10 ⁻⁹	2			>10 ⁻⁵

46	COMPOUND	ISO- MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	D1 IC50 (uM)	D2 IC50 (uM) (n)	PCP IC50 (M)
45			6.95×10^{-9}	7.50×10^{-10}	2			$>10^{-5}$
47	 CF ₃		5.65×10^{-9}	1.02×10^{-9}	2			$>10^{-5}$
48			2.70×10^{-9}	2.00×10^{-10}	2			$>10^{-5}$
49			1.038×10^{-8}	2.61×10^{-9}	4	>10 (2)	5.77 $\pm .83$ (3)	$>10^{-5}$
50			2.95×10^{-9}	5×10^{-11}	2			$>10^{-5}$

-71-

#	COMPOUND	ISO- MER	SIGMA IC50 (M)	SIGMA S.E.M.	n	D1 IC50	D2 (uM)	PCP IC50 (M)
51			4.23×10^{-9}	1.42×10^{-9}	4	2.99 $\pm .04$ (3)	.158 $\pm .028$ (2)	$>10^{-5}$

Results

As can be seen in Table 4, the sigma receptor ligands of the present invention exhibit very high binding with respect to the sigma receptors and very low binding with respect to the PCP and DA receptors. Therefore, these sigma receptor ligands can be used for the treatment of mental illness without the extrapyramidal side effects of traditional neuroleptic agents caused by binding to the DA receptor.

EXAMPLE 3 5-HT_{1A} BINDING ASSAYS

The sigma receptor ligands listed above (nos. 1-51) were further tested for binding at the 5-HT_{1A} receptor using the method of Peroutka, S.J., J. Neurochem. 47:529-540 (1986). The results appear in Table 5.

-73-

Table 5

Compound No.		5-HT _{1A} IC ₅₀ (nM)	\pm SEM(n)	Ratio 5-HT _{1A} / σ
5	1	115.00	12.00(2)	4
	2	90.30	28.83(3)	12
	3	1776.33	579.30(3)	.20
	4	4052.50	372.50(2)	5.9
	5	1500.00	100.00(2)	.88
10	6	3781.50	3458.50(2)	31
	7	2475.00	95.00(2)	40
	8	43.65	1.85(2)	.92
	9	122.00	6.00(2)	2
	10	212.00	31.00(2)	22
15	11	104.30	16.70(2)	11
	12	131.58	39.31(4)	56
	13	288.00	62.00(2)	2
	14	154.50	35.50(2)	5.4
	15	30.70	3.43(3)	2
20	16	44.8	2.5(2)	2.4
	17	87.80	12.20(2)	13
	18	588.50	68.50(2)	54
	19	114.23	12.43(3)	7.6
	20	108.05	10.95(2)	17
25	21	195.50	14.50(2)	21
	22	202.00	47.00(2)	13
	23	105.75	13.25(2)	25
	24	955.50	33.50(2)	1.4
	25	753.00	135.00(2)	7.3
30	26	10000	(1)	3.5
	27	6514.00	874.00(2)	200
	28	2442.50	342.50(2)	.05
	29	15515.00	5515.00(2)	3.1
	30	54999.50	44999.50(2)	2.6
35	31	19.30	4.20(2)	.92
	32	8.73	0.46(2)	.80
	33	51.60	2.30(2)	2.6
	34	6650.00	710.00(2)	2.6
	35	10360.00	360.00(2)	40
40	36	1626.67	378.17(3)	239
	37	20.30	4.70(2)	2.6
	38	107.75	27.25(2)	25
	39	89.60	0.10(2)	75
	40	2965.00	585.00(2)	970
45	41	34.67	9.09(3)	7.6
	42	572.50	134.50(2)	110
	43	1112.33	162.85*3)	50
	44	277.67	33.52(3)	40
	45	594.00	67.68(3)	100
50	46	21.47	3.60(3)	3

Table 5 (Cont.)





<u>Compound No.</u>	<u>IC₅₀ (nM)</u>	<u>5-HT_{1A} ±SEM(n)</u>	<u>Ratio 5-HT_{1A}/σ</u>
47	9.23	1.58(2)	1.6
48	6.30	0.74(2)	2.3
49	312.67	27.38(3)	30
50	281.67	69.00(3)	94

EXAMPLE 4 FURTHER [³H]DTG AND 5HT-1A BINDING ASSAYS

Further sigma receptor binding assays against [³H]DTG and 5HT-1A binding assays were conducted according to the procedure outlined in the previous examples. The results of these further studies are listed in Table 6.






-75-

TABLE 6

#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
52		11891.00	399.00(2)	--	--
53		10901.50	1198.50(2)	--	--
54		11815.00	1815.00(2)	--	--
55		6393.67	4426.77(3)	--	--

-76-

TABLE 6 (Cont.)

#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
5					
10					
15					
20					
25					
30					
35					
40					
45					
56		1190.50	159.50(2)	--	
57		6069.00	251.00(2)	--	
58		2054.50	14.50(2)	--	
59		2830.00	470.00(2)	--	
60		575.50	98.50(2)	--	

-77-

TABLE 6 (Cont.)










#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
61		14850.00	4850.00(2)	--	
62		11625.00	1625.00(2)	--	
63		763.33	352.86(3)	--	
64		77.40	4.29(3)	784.00	17.00(2)
65		72.87	3.46(3)	182.00	30.00(2)

TABLE 6 (Cont.)

#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
5					
10					
15		45.10	6.85(3)	272.00	44.00(2)
20					
25					
30		3.25	0.85(4)	1320.00	158.46(3)
35					
40		2.25	1.00(4)	891.00	181.56(3)
45					
50		116.20	62.80(2)	3885.75	1466.82(4)

-79-

TABLE 6 (Cont.)











#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
5					
10					
15	71 	5.68	2.19(2)	225.33	57.50(3)
20					
25	72 	7.23	1.92(2)	11.30	3.25(3)
30					
35	73 	6.98	2.16(2)	55.70	0.50(2)
40					
45	74 	81.87	21.83(3)	--	
50					
	75 	13.57	1.67(3)	255.00	(1)

TABLE 6 (Cont.)

#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
5					
10					
15					
20					
25					
30					
35					
40					
45					
76		2.65	0.63(2)	119.00	(1)
77		2.07	0.50(3)	759.00	(1)
78		3.01	0.23(2)	98.40	(1)
79		2.25	1.14(3)	173.00	(1)
80		130.10	18.16(3)	--	

-81-

TABLE 6 (Cont.)






#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
5					
10		0.88	0.18 (9)	43.90	(1)
15					
20		0.96	0.21 (9)	136.50	(1)
25					
30		7.08	3.93(2)	77.70	(1)
35					
40		6.43	0.50(2)	168.00	(1)
45					
		28.15	4.65(2)	1150.00	(1)

TABLE 6 (Cont.)






#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
86		2.43	1.12(2)	502.00	(1)
87		21.83	2.86(3)	67.40	(1)
88		53.40	9.80(3)	--	
89		2.74	0.40(4)	106.00	7.00(2)
90		3.95	0.53(4)	3208.00	148.00(2)

TABLE 6 (Cont.)






#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
91		4.07	0.52(4)	732.50	111.50(2)
92		2.99	0.28(4)	26.30	1.10(2)
93		1.40	0.05(4)	596.50	2.50(2)
94		4.02	0.27(4)	5549.50	119.50(2)
95		16950.00	(1)	--	--

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
96		141.00	29.61(3)	--	--
97		20.33	4.73(3)	--	--
98		6.18	1.12(4)	2.23	0.84(2)

TABLE 6 (Cont.)

#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)

5


10

100

25.65 0.75(2)

--

15



20


25

101

15.70 1.20(2)

--

30



35

40

102

6693.50 1423.50(2)

--




TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
103		75.20	36.80(2)	--	--
104		76.15	17.05(2)	--	--
105		9.50	3.38(3)	4956.50	565.50(2)

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
106		49.75	8.65(2)	--	--
107		10,000.00	0.00(2)	--	--
108		37.85	0.65(2)	--	--

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
109		184.50	12.50(2)	--	--
110		3.62	0.84(3)	116.90	22.20(2)
111		2.73	0.58(3)	92.20	22.10(2)

TABLE 6 (Cont.)







#	COMPOUND	SIGMA MEAN	IC ₅₀ SEM	(nM) (n)	5HT-1A MEAN	(nM) SEM(n)
112		2.65	0.41	(3)	8243.00	685.00(2)
113		S(+)	2.22	0.32	(3)	2148.00 105.00(2)
114		S(+)	6.71	1.07	(3)	34000 24,000(2)

TABLE 6 (Cont.)

#	COMPOUND	SIGMA		IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM	(n)		MEAN	SEM(n)
115		S(+)	3.04	0.42(3)		20000	10000(2)
116		S(+)	10.13	8.28(2)		--	
117			2.11	0.14(3)		781.50	258.50(2)

-91-

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM) MEAN SEM (n)	5HT-1A (nM) MEAN SEM(n)
5	<hr/>		
10		2.14 0.21(3)	1967.00 394.00(2)
15			
20			
25		21.40 4.80(2)	--
30			
35			
40		64.70 2.40(2)	--

TABLE 6 (Cont.)





#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
124		2.05	0.25 (3)	--	
125		38.83	9.05(2)	--	
126		219.33	7.45(3)	--	

TABLE 6 (Cont.)

#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)


5

10

127		10.57	1.28(3)	--
-----	---	-------	---------	----

15


20

128		2.87	0.72(3)	--
-----	--	------	---------	----

30

35

40

129		2.29	1.37(3)	--
-----	---	------	---------	----

45

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
5					
10					
15	130	4.23	0.14(3)	--	
20					
25					
30	131	46.90	15.45(3)	--	
35					
40					
45	133	S(+)	6.44	1.77(3)	--
					

TABLE 6 (Cont.)


#	COMPOUND	SIGMA MEAN	IC ₅₀ SEM (n)	(nM)	5HT-1A (nM) MEAN	SEM(n)
---	----------	---------------	-----------------------------	------	---------------------	--------

5

10

134 S(+) 198.77 74.48(3) --

15




20

25

135 13.50 4.37(3) --

30



35

40

136 1.78 1.32(3) --

45




TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
137		1.11	0.52(3)	--	
138		2.67	0.37(3)	--	
140		69.03	13.28(4)	--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
144		4.39	0.95(3)	--	
145		1.63	0.39(3)	--	
146		2.38	0.27(3)	--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
147		117.3	63.05(3)	--	
148		6.43	0.85(3)	--	
149		21.48	2.24(4)	--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA MEAN	IC ₅₀ SEM (n)	(nM)	5HT-1A MEAN	(nM) SEM(n)
5						
10						
15	150 	0.96	0.24(3)		--	
20						
25						
30	151 	1.39	0.10(3)		--	
35						
40						
45	152 	1.76	0.23(3)		--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
153		2.21	0.39(3)	--	
154		3955	711(3)	--	
155		1866	113(3)	--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
156		130.63	21.39(3)	--	--
157		6113	2790(2)	--	--
159		1.30	0.27(3)	--	--

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
160		0.62	0.14(3)	--	--
161		10.87	1.70(3)	--	--
162		0.90	0.05(3)	--	--

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
163		1.74	0.06(3)	--	
164		5.19	1.03(2)	--	
165		1.27	0.05(2)	--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA MEAN	IC ₅₀ (nM) SEM (n)	5HT-1A (nM) MEAN SEM(n)
166		1.03	0.01(2)	--
167		1.96	0.52(3)	--
169		22.80	2.30(2)	--

TABLE 6 (Cont.)




#	COMPOUND	SIGMA		IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM	(n)		MEAN	SEM(n)
170		4.18	0.56	(2)		--	
171		1.19	0.24	(2)		--	
172		7.46	1.03	(2)		--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
5					
10					
15		17.20	4.60(2)	--	
20					
25		1.10	0.35(3)	--	
30					
35					
40					
45		42.90	1.64(3)	--	

TABLE 6 (Cont.)




#	COMPOUND	SIGMA MEAN	IC ₅₀ (nM) SEM (n)	5HT-1A (nM) MEAN SEM(n)
177		7.37	1.57(2)	--
178		0.896	0.077(2)	--
179		2.82	0.02(2)	--

TABLE 6 (Cont.)




#	COMPOUND	SIGMA MEAN	IC ₅₀ SEM (n)	(nM)	5HT-1A MEAN	(nM)	SEM(n)
180		2.24	0.62	(2)	--		
181		1.08	0.06	(2)	--		
182		4.68	0.14	(2)	--		

TABLE 6 (Cont.)




#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
183		1.97	0.05(3)	--	--
184		1.98	0.14(2)	--	--
185		1736	732(2)	--	--

TABLE 6 (Cont.)







#	COMPOUND	SIGMA IC ₅₀ (nM)		5HT-1A (nM)	
		MEAN	SEM (n)	MEAN	SEM(n)
186		656	165(2)	--	
187		50.6	11.7(2)	--	
188		174	27(2)	--	

TABLE 6 (Cont.)

#	COMPOUND	SIGMA IC ₅₀ (nM) MEAN SEM (n)	5HT-1A (nM) MEAN SEM(n)
5			
10			
189		13.5 3.3(2)	--
15			
20			
25			
190		1.07 0.07(2)	--
30			
35			
40			
191		8.14 -- (1)	--
45			
50	<u>EXAMPLE 5</u> <u>STRUCTURE-ACTIVITY RELATIONSHIP OF THE SIGMA RECEPTOR LIGANDS</u>		

The binding data reproduced in Tables 4-6 allows the identification of groups of compounds having high binding and

selectivity for the sigma receptors. For example, it has been determined that the α -methyl group on N-(3-phenylpropyl)isopropylamine ($K_i = 22$ nM) is not required for high binding to the sigma receptor. The corresponding des-methyl compound N-(3-phenylpropyl)-2-phenethylamine binds with comparable affinity ($K_i = 19$ nM). This des-methyl compound possesses two carbon atoms between the phenyl ring and the terminal amine and, on the other side of the molecule, possesses three carbon atoms between the amine and the second phenyl ring (see Table 7; $x = 2$, $y = 3$). A systematic comparison of phenylalkylamine derivatives with varying x and y values is shown in Table 7.

As shown in Table 7, the increase of x from 2 to 3 resulted in the retention of affinity. Interestingly, the total carbon chain length does not appear to be critical for high binding to the sigma receptor. The highest affinity agents are those where either x or $y = 5$. Thus, a phenylpentylamine is optimal while the length of the chain on the other side of the molecule appears to be inconsequential.

Since a carbon length of five atoms is optimal for high binding, and if the substituents on the other side of the molecule are relatively less important, then the second phenyl ring may not be important. Therefore, N-methyl-N-propyl-5-phenylpentylamine, a des-phenyl analog, was prepared and found to bind with high affinity ($K_i = 2.2$ nM).

Where x and $y = 3$, it has also been found that replacement of one of the phenyl rings with a cyclohexyl ring ($K_i = 2.5$ nM) resulted in a nearly six-fold increase in affinity. Furthermore, the cyclohexyl analog N-benzyl-(5-cyclohexyl)pentylamine, where the cyclohexyl group is separated from the amine by five carbon atoms, also binds with high affinity ($K_i = 1.3$ nM).

Taken together, these data suggest that (a) phenylpentylamines bind with high affinity to the sigma receptor, (b) that the nature of the amine substituent is relatively unimportant, and (c) that the phenyl ring of the phenylpentylamines may be replaced by a cyclohexyl ring with retention (and an increase) of affinity.

Table 7
 Phenyl-(CH₂)_x-NH-(CH₂)_y-Phenyl

	x	y	Ki (nM)
5	1	1	175
	1	4	13
10	1	5	2.0
	1	7	3.6
	2*	3	19
	2	4	5.5
15	2	5	2.4
	2	7	4.1
	3	3	14
	3	4	7.0
20	3	5	1.8
	3	6	2.6
	4	4	6.7
	4	5	2.5
25	5	1	2.0
	5	2	2.4
	5	3	1.8
	5	4	2.5


* = des-methyl N-(3-phenylpropyl)isopropylphenylamine

With regard to sigma selectivity over 5-HT_{1A}, the unsubstituted phenyl derivative N-(3-phenylpropyl)-1-isopropylamine (cmp. no. 1) exhibits only a 4-fold selectivity for sigma receptors. Aromatic substitution enhances affinity for sigma receptors whereas 5-HT_{1A} affinity remains relatively constant. Consequently, these aromatic substituted derivatives bind with a relatively low, but constant selectivity. Next, the effect of terminal amine modification was examined where the aromatic portion was held constant as a phenyl group. Replacement of the benzylic methylene group of cmp. #1 by an oxygen atom, carbonyl group, or sp-hybridized carbon atom (cmps. 8, 9 and 25) gave little effect. Removal of the α -methyl group of cmp. #1 (R(-)) seems to enhance 5-HT_{1A} affinity. The optical

isomers of cmp. #1 bind at sigma receptors with comparable affinity; however, the S(+) isomer binds at 5-HT_{1A} receptors with only one-tenth the affinity of its antipode resulting in a 48-fold selectivity. Similar results were obtained with the isomers of N-(2-ethylphenyl)isopropylamine (cmp. #7). N-Monomethylation (e.g. cmps. 45 and 69) also seems to enhance sigma affinity and selectivity. To further substantiate this finding, N-methyl-N-(3-propylphenyl)-1-(4-n-propylphenyl)-isopropylamine was prepared and found to have the highest and most selective affinity for the sigma receptor.

EXAMPLE 6 Further Structure-Activity Studies of 5-Pentylamine Derivatives

With N-substituted 5-phenylpentylamines, the length of alkyl chain that separates the amine from its aromatic (phenyl-B) substituent has little influence on affinity (e.g., compare 76, 77, 89, and 111; K_i = 2.0-2.7 nM). It seems unlikely that the phenylethylamines and phenylpentyl amines bind in exactly the same manner at the sigma receptors due to the difference in chain length. In order to account for the binding of these compounds, the phenylethylamines and the phenylpentylamines may utilize different aromatic binding sites. These results prompted a further investigation of 5-phenylpentylamine derivatives as sigma ligands.



76	n = 1
77	n = 2
89	n = 3
111	n = 4

CHEMISTRY: The compounds in Table 10 were synthesized by one of the methods described (A-H). Most of the compounds were prepared in two or three steps using either acylation and reduction of the

intermediate amide or direct alkylation procedures. The amides were prepared either from the acyl halide and an amine or directly using ethyl chloroformate, an appropriate acid and an amine. Subsequent reduction by lithium aluminum hydride afforded the target amines. Following this procedure, the benzyl protected analog of compound 135 was obtained and subsequent hydrogenolysis produced the desired compound. Compounds 136 and 152 were obtained by methylating the corresponding secondary amines using Eschweiler-Clark reductive alkylation procedure. Compounds 173 and 172 were prepared by O-demethylation of 144 and 145 respectively, using concentrated hydrobromic acid solution. Other target compounds prepared either by direct alkylation (method C) or by reductive alkylation (method E), using suitable aldehydes and appropriate amines.

EXPERIMENTAL:

Synthesis: Proton magnetic resonance spectra were obtained on JEOL FX90Q or QE 300 (300 MHz) spectrometers with tetramethylsilane as internal standard. All spectra are consistent with the assigned structures. Melting points were determined on a Thomas Hoover apparatus and are uncorrected. Elemental analyses were performed by Atlantic Microlab and determined values are within 0.4% of calculated values.

Method A:

N-Cyclohexylmethyl-5-cyclohexylpentylamine Hydrochloride (151). A solution of ethyl chloroformate (1.8g, 8.2 mmol) in methylene chloride (25 mL) was added in a dropwise manner to a stirred ice-cooled solution of cyclohexanepentanoic acid (3g, 8.1 mmol) and Et₃N (1.7g, 8.1 mmol) in dry methylene chloride (50 mL) under N₂, over 10 min. Stirring was continued for 30 min. and cyclohexylmethylamine (1.8g, 8.1 mmol) in methylene chloride (25 mL) was added dropwise over 5 mins. Stirring was allowed to continue for an additional 3 h after which the reaction mixture was washed water (50 mL) and dried (MgSO₄). Solvent was removed

in vacuo to afford an oil which was crystallized from MeOH-H₂O (4.1g, 91%); mp 69-72°C. A solution of the amide (3.8g, 14 mmol) in THF (50mL) was added dropwise to a suspension of LiAlH₄ (2.6g, 5 eq) in THF (100 mL). The reaction mixture was heated under reflux in a stream of N₂ for 20 h. Excess LiAlH₄ was decomposed by the gentle addition of H₂O and 10% NaOH solution. Solid matter was removed by filtration and solvent was removed in vacuo to obtain an oily residue. The residue was taken up in Et₂O, dried (MgSO₄) and solvent was removed under reduced pressure to afford an oil (3.0 g, 80%). The hydrochloride salt, obtained by the addition saturated solution of ethereal HCl to an ethereal solution of 151 (free base), was recrystallized MeOH/EtOAc; mp 223-224°C (see Table 8).

Method B:

N-Methyl-N-hexyl-2-phenylethylamine Hydrogen Oxalate (130).

A solution of hexanoyl chloride (1g, 7.4 mmol) in THF (40 mL) was added in a dropwise manner to a stirred solution of N-methyl-2-phenylethylamine (1g, 7.4 mmol) and Et₃N (2.3 g, 20 mmol) in THF (100 mL) cooled to 0°C. The reaction mixture was allowed to stir overnight (20 h) after which the triethylamine salt was removed by filtration and washed with THF (2 x 20 mL). The combined filtrate and washings were evaporated under reduced pressure and the residue taken up in CHCl₃ (30 mL). The chloroform solution was washed with H₂O (30 mL) and dried (Na₂SO₄). Solvent was removed in vacuo to obtain an oil which was shown by IR (C=O, 1644cm⁻¹) to be an amide (1.5g, 87%). A solution of the amide (0.5g, 2.1 mmol) in dry THF (60 mL) was added dropwise to a suspension of LiAlH₄ (0.41g, 11 mmol) in THF(40 mL) under a stream of N₂. The mixture was stirred overnight, cooled to 0°C and the reaction mixture was quenched by cautious addition of water and 15% NaOH solution. Solids were removed by filtration. The filtrate was evaporated to dryness under reduced pressure and the residue was taken up in Et₂O (50 mL and dried (MgSO₄). Solvent was removed under reduced pressure to obtain an oil (350 mg, 75%). The oil was converted to the oxalate salt (white

crystals) and recrystallized from MeOH/Et₂O solution; mp 139-140°C (see Table 8).

Method C:

5 N-Methyl-N-propylhexylamine Hydrogen Oxalate (131). A stirred mixture of N-methylpropylamine (1.2g, 13.7 mmol), 1-bromohexane (3.4g, 20.5 mmol) and potassium carbonate (3.8g, 27 mmol) in 1,2-dimethoxyethane (DME) (8 mL) was heated under reflux for 24 h and allowed to cool to room temperature. The solid material was removed by filtration and washed several times with CHCl₃. The combined filtrates were evaporated to dryness under reduced pressure and the residue partitioned between Et₂O (30 mL) and 10% NaOH solution (20 mL). The ethereal fraction was washed with H₂O (10 mL) and dried (Na₂SO₄). A saturated solution of oxalic acid in anhydrous Et₂O was added to obtain a white solid which was recrystallized from iPrOH/Et₂O (0.4g, 12%); mp 101-102°C (see Table 8).

Method D:

20 N-Methyl-N-cyclohexylmethyl-5-cyclohexylpentylamine Hydrochloride (152). A mixture of 151 [CNS#?] (0.5g, 1.9 mmol), formic acid (1.1g, 23 mmol) and formaldehyde solution (37%) (1.85g, 23 mmol) was heated at about 100°C for 22 h and allowed to cool to room temperature. A 3N HCl solution (10 mL) was added and the solution extracted with Et₂O (3 x 25 mL). The ethereal solution, which contained the expected product, was washed with 10% NaOH (30 mL), then water (10 mL) and dried (MgSO₄). A saturated ethereal HCl solution was added, solvent removed in vacuo, and the residue recrystallized from MeOH-EtOAc (370mg, 61%); mp 162-163°C (see Table 8).

Method E:

35 N-(3-Cyclohexylpropyl)-3-phenylpropylamine Hydrochloride (92). A solution of 3-phenyl propylamine (0.65 g, 5 mmol) and 3-cyclohexylpropionaldehyde (0.75 g, 5.4 mmol) in MeOH (40 mL) was hydrogenated in a Parr bottle containing 10% Pd/C (0.3 g) until

sufficient H_2 was taken up (40 min). The catalyst was removed by filtration; the filtrate was concentrated to about 10 mL under reduced pressure and added to 1N HCl solution (20 mL). The precipitate was collected by filtration and washed thoroughly with anhydrous Et_2O (3 x 10 mL). Recrystallization from 2-butanone afforded the desired compound as white shiny plates (0.75 g, 53%); mp 203-205°C (see Table 8).

Method F:

N-[2-(3-hydroxyphenyl)ethyl]-5-phenylpentylamine Hydrobromide (172). A mixture of 145 (free base) (0.19 g, 0.65 mmol) and hydrobromic acid solution (48%) (0.22 mL, 1.3 mmol) was heated at reflux for 2 h and solvent was removed in vacuo. The solid residue was recrystallized from MeOH/anhydrous Et_2O to afford the desired compound (100 mg, 42%); mp 151-153°C (see Table 8).

Method G:

N-Phenyl-5-phenylpentylamine Hydrogen Oxalate (135). A mixture of N-benzyl-N-phenyl-5-phenylpentylamine (0.89 g, 2.5 mmol) in EtOH (20 mL) and 10% Pd/C (0.1 g) was hydrogenated at 50 psi for about 3 h. The catalyst was removed by filtration and the filtrate was evaporated to dryness under reduced pressure. The residue was partitioned between 10% HCl solution (20 mL) and Et_2O (20 mL); the aqueous portion was basified with 10% NaOH and extracted with Et_2O (20 mL). The Et_2O portion was dried ($MgSO_4$) and solvent was removed in vacuo to afford an oil (0.3 g, 49%). The oxalate salt was prepared by the addition of a saturated solution of oxalic acid and subsequently recrystallized from EtOAc; mp 133-134°C (see Table 8).

Method H:

5-Cyclohexylpentylamine Hydrogen Oxalate (126). A solution of cyclohexanepentanoic acid (2 g, 10.9 mmol) in $SOCl_2$ (8 mL) was heated on a steam bath for 3 h. The $SOCl_2$ was removed under vacuum. Chloroform (2 x 10 mL) was added and reevaporated to give the crude acid chloride as a yellow liquid. A saturated

solution of ammonia in dry THF (ammonia gas bubbled through 10 mL of THF for 5 min), was slowly added to a solution of the acid chloride in dry THF (50 mL) while cooling in an ice bath. The reaction mixture was allowed to stir at room temperature for 3 h and was then concentrated to half of its volume and poured onto water (100 mL). The solid was removed by filtration, washed with water and dried (MgSO_4). Recrystallization from aqueous MeOH gave 5-cyclohexylvaleramide (1.1g, 55%); mp 121-122°C (reported mp 122-123°C) (Katsellson and Dubinin, *Compt. Rend. Acad. Sci. U.R.S.S. [N.S.]* 4:405 (1936); *Chem. Abstr.* 31:3449 (1937)). A suspension of the amide (183 mg, 1 mmol) in anhydrous Et_2O (9 mL) was slowly added to a stirred suspension of LiAlH_4 (114 mg, 3 mmol) in Et_2O (15 mL). After addition was complete, the mixture was heated at reflux for 2 h and then allowed to stir overnight at room temperature. The reaction mixture was cooled to 0°C and excess LiAlH_4 was decomposed by successive addition of water (2 mL), 2N NaOH (2 mL) and water (5 mL). After stirring vigorously for 20 min, the mixture was filtered and the inorganic residue was washed with warm Et_2O (3 x 8 mL). The combined Et_2O solution was dried over anhydrous K_2CO_3 and evaporated to dryness. The residue was distilled under reduced pressure to yield a clear oil (99 mg, 59%); bp 30-36°C (0.35 mm), Lit. bp 108-113°C (15 mm) (Skinner, C.G., et al., *J. Am. Chem. Soc.* 79:2844 (1957)). A solution of the free base in anhydrous Et_2O was added to a solution of oxalic acid (52 mg) in anhydrous Et_2O (8 mL) with continuous shaking. The precipitated solid was recrystallized from MeOH/anhydrous Et_2O ; mp 164°C (see Table 8).

Radioligand Binding. The σ binding assay was conducted using guinea pig (Taconic) brain membranes and [^3H]di-o-tolylguanidine (DTG) as radioligand. Briefly, membranes (P2 microsomal fraction) were diluted 1:3 with 50 mM Tris HCl (pH 7.4) and 0.4 mL was combined with 50 μL [^3H]DTG (1-2 nM final concentration) and 50 μL of competing drug or buffer. After 90 min at room temperature, incubation was terminated by rapid filtration under vacuum through Whatman GF/B glass fiber filters using a Brandel

48-well cell harvester. Filters were washed three times with 5 mL of cold Tris HCl buffer and each filter was suspended in 5 mL of Cytoscint (ICN Biomedical). Radioactivity was measured by liquid scintillation spectrometry at a counting efficiency of 50%. Non-specific binding was measured in the presence of 10 μ M haloperidol.

Microanalytical Data Calculated/Found				
	C	H	N	
10	<u>152</u>	70.56	7.61	3.92
		70.43	7.59	3.93
	<u>136</u>	71.66	8.11	3.63
		71.37	8.08	3.59
15	<u>144</u>	71.94	8.45	4.20
		71.88	8.49	4.19
	<u>8</u>	67.54	7.29	3.75
		67.47	7.33	3.69
20	<u>145</u>	71.94	8.45	4.20
		71.87	8.49	4.21
	<u>172</u>	62.02	7.12	3.81
		62.01	7.22	3.81
25	<u>146</u>	71.18	8.48	4.15
		71.23	8.46	4.17
	<u>135</u>	68.90	7.06	4.23
		68.52	7.22	4.23
30	<u>112</u>	65.99	8.80	4.53
		65.82	8.75	4.51
	<u>130</u>	65.99	8.80	4.53
		65.92	8.79	4.57
30	<u>131</u>	58.27	10.19	5.66
		58.20	10.17	5.66
	<u>193</u>	73.06	10.22	4.75

		72.95	10.23	4.74
92		73.06	10.22	4.73
		72.96	10.22	4.77
126		58.10	9.76	5.23
		58.18	9.77	5.22
127		61.60	9.97	5.12
		61.51	9.96	5.12
128		62.71	10.16	4.92
		62.69	10.17	4.87
125		63.94	8.19	5.10
		64.03	8.24	4.98
94		73.06	10.22	4.73
		73.12	10.27	4.78
23		70.55	12.01	4.57
		70.36	12.01	4.57
24		71.21	12.11	4.37
		71.27	12.04	4.33

TABLE 8. Physicochemical properties of new sigma receptor ligands

Compd #	Method	Recryst. Solv.	Mp (°C)	% Yield ^a	Formula
152	A	PrOH/EtOAc	140-141	49	C ₁₉ H ₂₅ N.C ₂ H ₂ O ₄
135	D	PrOH	141-142	73	C ₂₁ H ₂₉ N.C ₂ H ₂ O ₄
144	A	MEK	115	37	C ₂₀ H ₂₇ NO.HCl
173	B	MEK	173-174	2	C ₁₉ H ₂₅ NO.C ₂ H ₂ O ₄
145	A	MeOH/Et ₂ O	160	40	C ₂₀ H ₂₇ NO.HCl
172	F	MeOH/Et ₂ O	151-153	39	C ₁₉ H ₂₅ NO.HBr ^b
146	A	MeOH/Et ₂ O	233-235	33	C ₂₀ H ₂₇ NO.HCl ^b
135	G	EtOAc	133-134	35	C ₁₇ H ₂₁ N.C ₂ H ₂ O ₄ ^c
112	E	EtOAc	130-131	24	C ₁₅ H ₂₃ N.C ₂ H ₂ O ₄
130	B	MeOH/Et ₂ O	139-140	65	C ₁₅ H ₂₃ N.C ₂ H ₂ O ₄
131	C	iPrOH/Et ₂ O	101-102	12	C ₁₀ H ₂₃ N.C ₂ H ₂ O ₄
93	A	iPrOH/Et ₂ O	171-173	39	C ₁₅ H ₂₉ N.HCl

92	E	MEK	203-205	53	C ₁₈ H ₂₉ NHCl
126	H	MeOH/Et ₂ O	164	48	C ₁₁ H ₂₃ N.C ₂ H ₅ O ₄ ^d
127	B	MeOH	191	28	C ₁₂ H ₂₅ N.C ₂ H ₅ O ₄
128	B	MeOH	152	25	C ₁₃ H ₂₇ N.C ₂ H ₅ O ₄
125	B	MeOH	163	43	C ₁₃ H ₂₁ N.C ₂ H ₅ O ₄
94	A	iPrOH/Et ₂ O	184-185	34	C ₁₈ H ₂₉ NHCl
151	A	MeOH/ EtOAc	223-224	73	C ₁₈ H ₃₅ NHCl ^e
152	D	MeOH/ EtOAc	162-163	61	C ₁₉ H ₃₇ NHCl ^e
(a) No attempt was made to optimize yields. Yields are overall % yields. (b) Crystallized with 0.2 moles of H ₂ O. (c) Crystallized with 0.1 moles of H ₂ O. (d) Crystallized with 0.5 moles of H ₂ O. (e) Crystallized with 0.25 moles of H ₂ O.					


RESULTS and DISCUSSION

Binding data are shown in Table 9. N-Methylation of N-substituted phenylethylamines typically doubles and aromatic substitution at the 3- and 4-position has essentially no effect on sigma receptor affinity. N-Methylation of 151 and 111 (i.e., 152 and 136, respectively) doubles affinity (Table 9). Incorporation of phenolic or methoxy groups at the 3- and 4-positions (145, 172, and 146) has little effect on affinity, whereas the 2-hydroxy derivative 8 binds with about 7-fold lower affinity than its unsubstituted parent 89 (K_i = 2.4 nM). Excision of the N-alkyl chain separating the amine from phenyl-B, to afford the aniline derivative 135 (K_i = 12 nM), reduces affinity by only 6-fold; this decrease in affinity may reflect a decrease in the basicity of the amine.

The above results (i.e., lack of influence of chain length and aromatic substitution on sigma affinity) suggest that the aromatic amine substituent plays only a small role in binding, and that it may be possible to replace the phenyl-B ring with a non-aromatic group. Indeed, compound 122, which may be viewed either as either an N-methyl analog of 89 where the phenyl-B group has been replaced by methyl, or as an analog of 136 where

the phenyl group has been eliminated, binds with high affinity ($K_i = 2.4$ nM). Interestingly, the phenyl-A ring can also be replaced by a methyl group (130; $K_i = 3.8$ nM) with little reduction in affinity. Replacement of both phenyl groups by methyl (131; $K_i = 45$ nM) results in a significant reduction in affinity. Apparently, either phenyl-A or phenyl-B can be replaced by a methyl group; however, replacement of both results in decreased affinity.

Because both phenyl groups are unnecessary for binding, we examined several additional compounds bearing a single phenyl. Compound 93, the phenyl-B reduced analog of 151, binds with twice the affinity ($K_i = 1.2$ nM) of 151. Compound 92 ($K_i = 2.4$ nM), an analog of 93 where the position of the amine has been shifted, also binds with high affinity.



Examination of three simple 5-(cyclohexyl)pentylamines, the primary amine 126 ($K_i = 200$ nM), the N-monomethylamine 127 (K_i 10 nM), and the N,N-dimethylamine 128 ($K_i = 2.6$ nM), reveals that tertiary amines are most favorable for binding. The affinity of the latter compound 128, was compared with the aromatic analog 125 ($K_i = 35$ nM), revealing that reduction results in about a 10-fold increase in affinity. However, reduction of the phenyl-A ring of 151 (94; $K_i = 3.4$ nM) results in essentially no change in affinity. This may be related to the fact that 128 and 125 are tertiary amines whereas 94 is a secondary amine; nevertheless, it is apparent from these results that both phenyl-A and phenyl-B can be replaced with a cyclohexyl group. The final two compounds examined, 151 and 152, possess two cyclohexyl shortened or lengthened without adverse effect on sigma receptor affinity (e.g., see compounds 151, 89, 111 and 76). That is, affinity is independent of the length of the alkyl chain that separates the

phenyl-B ring from the amine. To account for these observations, the sigma receptors may possess at last two distinct aromatic binding sites: one that utilizes the phenyl-B ring of phenylethylamines, and one that utilizes the phenyl-A ring of the phenylpentylamines. The phenyl-B ring of 151, 89, 111 and 76 is unnecessary for binding; it can be replaced by a small alkyl group (e.g. 112) or with a cyclohexyl group (e.g. 131) with retention of affinity. Interestingly, the phenyl-A ring can also be replaced with a methyl or cyclohexyl group (e.g., 130 and 94, respectively) with little change in affinity. Had only one of the phenyl groups been replaceable by methyl (or cyclohexyl) without loss of affinity, this would have provided direct evidence for the two distinct sites on sigma receptors. Nevertheless, in as much as compounds such as 151 and 152 bind with high affinity, it would appear that, independent of mode of binding, an aromatic moiety is not a requirement for high affinity.

Table 9 Sigma receptor binding data.*



	X	R	n	Y	Ki nM (SEM)
77	Phenyl	H	1	Phenyl	2.0
89	Phenyl	H	2	Phenyl	2.4
111	Phenyl	H	3	Phenyl	2.7
76	Phenyl	H	4	Phenyl	2.5
137	Phenyl	Me	1	Phenyl	1.0(±0.4)
136	Phenyl	Me	3	Phenyl	1.6(±1.2)
144	Phenyl	H	2	2-OMe Phenyl	4.0(±0.9)
173	Phenyl	H	2	2-OH Phenyl	15.6(±4.1)
145	Phenyl	H	2	3-OMe Phenyl	1.5(±0.4)
172	Phenyl	H	2	3-OH Phenyl	6.8(±0.9)
146	Phenyl	H	2	4-OMe Phenyl	2.2(±0.3)
135	Phenyl	H	0	Phenyl	12.2(±4.0)
112	Phenyl	Me	2	Methyl	2.4(±0.4)
130	Methyl	Me	2	Phenyl	3.8(±0.1)
131	Methyl	Me	2	Methyl	45(±14)
93	Phenyl	H	1	Cyclohexyl	1.2(±0.0)
126	Cyclohexyl	H	0	H	200(±7)
127	Cyclohexyl	H	0	Methyl	10(±1)
128	Cyclohexyl	Me	0	Methyl	2.6(±0.7)
125	Phenyl	Me	0	Methyl	35(±8)
94	Cyclohexyl	H	1	Phenyl	3.4(±0.3)
151	Cyclohexyl	H	1	Cyclohexyl	1.2(±0.1)
152	Cyclohexyl	Me	1	Cyclohexyl	1.6(±0.2)

EXAMPLE 7 Discrimination between the Sigma-1 and Sigma-2 Binding Sites

Materials

[³H](+)pentazocine (35 Ci/mmol) was generously provided by Dr. Steven Hunt of DuPont/NEN. [³H]DTG was purchased from DuPont/NEN. (+)Pentazocine was obtained from the NIDA Research Technology Branch, Division of Research. Guinea pig whole membrane preparations (P2 microsomal fraction) were obtained and used as described by Weber, E. et al., Proc.Natl.Acad.Sci 83:8784-8788 (1986).

Sigma-1 binding assay ([³H](+)pentazocine)

The sigma-1 selective binding assay was performed using [³H](+)pentazocine as the radioligand (3-4 nM final concentration unless otherwise specified) and approximately 100 µg of guinea pig membranes in a final volume of 500 µl of 50 mM TRIS-HCl, pH 8.0. Non-specific binding was determined in the presence of 10 µM haloperidol. For the standard equilibrium assay, the mixtures were incubated for 4-5 hours at 37°C, quenched with 4 ml of ice cold incubation buffer and rapidly filtered over Whatman GF/B fiber filters, followed by three 4 ml rinses with additional ice cold incubation buffer. The radioactivity on the filters was determined by scintillation spectrometry at an efficiency of about 50 % using Cytoscint (ICN) scintillation fluid.

Sigma-2 binding assay

The sigma-2 selective binding assay was performed using about 2 nM [³H]DTG as the radioligand in the presence of 200 nM (+)pentazocine to block the sigma-1 sites, with 400 µg of guinea pig membranes in a total volume of 0.5 ml of 50 mM TRIS-HCl, pH 7.4. Non-specific binding was determined in the presence of 10 µM haloperidol. For the standard equilibrium assay, the mixtures were incubated for 30 min. at room temperature, then filtered and the radioactivity determined as described above.

Data Analysis

Equilibrium binding data were analyzed by least squares nonlinear regression techniques as described by Fischer, J.B. and Schonbrunn, A., J. Biol. Chem. 263:2808-2816 (1988). Two site binding curves were fit to an equation describing the sum of two independent sites with Hill slopes of 1. Affinity constants (K_i) values were calculated from IC_{50} values using the Cheng-Prusoff equation (Cheng, Y.-C. and Prusoff, W.H., Biochem. Pharm. 22:3099-3108 (1973)). The results are reported in Table 10 for DTG, several lots of an N,N'-disubstituted guanidine, haloperidol and BMY-14802, as well as for certain of the compounds of the invention. For comparison, Table 11 shows the sigma-1 and sigma-2 binding data for a number of other known compounds.

TABLE 10

cap. #		SIGMA-1 K _i VS. 3H-PENTAZOCINE (N M)			SIGMA-2 K _i VS. 3H- DTG+PENTAZOCINE (N M)			S1/S2 Ratio
		MEAN	SEM	n	MEAN	SEM	n	
	N,N'-Di-(o-tolyl)guanidine	40.67	1.67	3	52.39	4.14	4	0.776
	Haloperidol	0.47	0.09	3	11.50	0.55	3	0.041
	BMY-14802	271.00	52.00	2	132.60	43.86	3	2.044
25	1 R(-) N-(1-phenyl-2-propyl)-3-phenylpropylamine	10.75	2.06	2	60.75	1.85	2	0.177
	10 (±) N-(1-phenyl-2-propyl)-4-phenylbutylamine	4.68	1.26	3	53.20	6.00	2	0.088
	11 R(-) N-(1-phenyl-2-propyl)-3-(1-naphthyl)propylamine	8.60	1.40	2	280.00	31.00	2	0.31
	12 R(-) N-(1-phenyl-2-propyl)-3-(2-naphthyl)propylamine	5.73	1.71	3	258.00	48.00	2	0.022
	15 Di-(3-phenylpropyl)-amine	11.37	2.00	3	63.90	1.37	3	.178
30	16 (±) N-[1-(1'-Naphthyl)-2-propyl]-3-phenylpropylamine	9.25	0.75	2	152.00	14.00	2	0.061
	17 (±) N-[1-(2'-Naphthyl)-2-propyl]-3-phenylpropylamine	31.00	13.00	2	222.00	82.00	2	0.140
	18 N-(3-phenylpropyl)-1-(2-naphthyl)-2-propylamine	25.70	6.30	2	15.13	2.61	3	1.7
	20 (±) N-[1-(3'-Bromophenyl)-2-propyl]-3-phenylpropylamine	9.65	2.35	2	26.15	10.35	2	0.369
	21 (±) N-[1-(4'-Bromophenyl)-2-propyl]-3-phenylpropylamine	11.97	2.85	3	39.13	9.20	4	0.306
35	22 (±) N-[1-(4'-Iodophenyl)-2-propyl]-3-phenylpropylamine	17.67	3.64	3	76.03	6.29	4	0.232

cmp. #		SIGMA-1 Ki vs. 3H-PENTAZOCINE			SIGMA-2 Ki vs. 3H- DTG-PENTAZOCINE			S1/S2 Ratio	
		MEAN	SEM	n	MEAN	SEM	n		
10	23	(±) N-[1-(3'-Trifluoromethyl)-2-propyl]-3-phenylpropylamine	8.75	4.05	4	20.85	3.48	4	0.420
	32	1-(3'-Chlorophenyl)-4-(2-phenylethyl)piperazine	14.95	7.40	4	42.43	4.02	3	0.352
	34	(±) 1-Phenyl-2-aminopentane	1177.00	415.64	3	10276.00	276.00	3	0.115
	40	4-Hydroxy-4-phenyl-1-(3-phenylpropyl)piperidine	0.68	0.36	3	9.20	2.07	3	0.074
	40	4-Hydroxy-4-phenyl-1-(3-phenylpropyl)piperidine	0.68	0.36	3	7.03	2.87	3	0.096
	43	N-(3-phenylpropyl)-1-phenyl-2-propylamine	51.47	11.44	3	37.50	11.02	3	1.37
	44	N-(4-phenylbutyl)-1-phenyl-2-propylamine	19.43	4.96	3	35.80	3.34	3	.543
	45	N-(5-phenylpentyl)-1-phenyl-2-propylamine	0.88	0.21	4	18.47	2.27	3	.048
	60	N-(2-Naphthyl)piperazine	219.87	60.75	3	5599.00	610.04	3	0.039
	64	N-Phenyl-N'-propylpiperazine	81.60	13.29	3	538.00	42.00	3	0.152
15	70	Di-[3-(2'-Naphthyl)propyl]amine	--	--	0	10000.00	0.00	2	--
	71	N-(4-phenylbutyl)phenethylamine	2.60	0.31	4	118.67	14.99	3	0.022
	74	Di-N-[3-(2'-naphthyl)propyl]-N-methylamine	29.98	11.06	3	460.75	165.40	4	0.065
	75	N-(4-phenylbutyl)benzylamine	9.61	2.88	5	161.63	31.37	3	0.059
	76	N-(5-phenylpentyl)-(4-phenyl)butylamine	0.48	0.05	2	68.00	5.00	2	0.007
	77	N-(5-phenylpentyl)benzylamine	0.32	0.10	3	34.33	6.87	3	0.009
	79	N-(4-phenylbutyl)-N'-benzylpiperazine	0.17	0.02	4	6.99	1.14	3	0.024
	80	N-4(4-phenylbutyl)-N'-benzoylpiperazine	71.83	2.51	3	965.00	65.00	2	0.074
	83	N-(3-phenylpropyl)-1-(p-ethoxyphenyl)-2-propylamine	3.38	0.30	2	33.50	0.50	2	0.101
	87	N-(3-phenylpropyl)phenethylamine	11.27	0.97	2	90.10	37.90	2	0.125
20	89	N-(5-phenylpentyl)phenethylamine	0.17	0.00	2	14.60	5.31	2	0.012
	90	N-(7-phenylheptyl)benzylamine	2.30	1.00	2	38.47	9.44	3	0.060
	91	N-(7-phenylheptyl)phenethylamine	1.47	0.63	4	33.23	10.16	3	0.044
	94	N-(5-cyclohexylpentyl)benzylamine	0.81	0.17	3	43.57	6.76	3	0.019
	96	N-phenyl-N'-(3-(1-phthalimido)propyl)piperazine	3100.33	1597.63	3	149.18	18.06	4	87.819
	97	N-phenyl-N'-(4-(1-phthalimido)butyl)piperazine	195.33	31.78	3	20.38	0.88	4	9.587
	101	N-(4-phenylbutyl)-1-phenyl-2-butylamine	2.70	0.60	2	104.10	12.90	2	0.026
	103	N-(4-phenyl-3(E)-butenyl)-N'-methylpiperazine	28.67	3.67	3	2544.67	699.93	3	0.011

		SIGMA-1 K1 vs. 3H-PENTAZOCINE			SIGMA-2 K1 vs. 3H-DTG+PENTAZOCINE			S1/S2 Ratio	
cmp.		MEAN	SEM	n	MEAN	SEM	n		
5	104	N-(4-phenyl-3(Z)-butenyl)-N'-methylpiperazine	28.67	6.67	3	1306.00	54.00	2	0.022
	105	N-(4-(3-trifluoromethyl)-3-(Z)-butenyl)-N-methylpiperazine	1.50	0.26	3	176.53	53.26	3	0.008
	106	N-(4-phenyl)-N'-methylpiperazine	19.33	3.93	3	898.80	180.96	5	0.022
	111	N-(5-phenylpentyl)-3-phenylpropylamine	0.28	0.03	3	9.84	4.58	3	0.029
	112	N-methyl-N-propyl-5-phenylpentylamine	0.29	0.05	3	40.48	10.77	5	0.007
	113	N-Methyl-N-(3-phenylpropyl)-1-phenylisopropylamine	0.51	--	1	4.87	0.72	3	0.105
10	117	N-(5-phenylpentyl)-4-benzylpiperidine	0.58	0.26	3	2.78	0.33	3	0.210
	120	N-(4-(1-phthalimido)butyl)-N'-(o-methoxyphenyl)piperazine	511.67	115.79	3	76.57	24.14	3	6.683
	124	N-(5-phenylpentyl)piperidine	0.48	0.01	3	49.90	8.87	.	0.010
	125	N,N-Dimethyl-5-phenylpentylamine	10.45	1.07	3	965.00	55.00	2	0.011
	126	5-cyclohexylpentylamine	191.33	16.76	3	2554.00	64.00	2	0.075
	127	N-methyl-5-cyclohexylpentylamine	6.78	0.92	4	346.00	29.00	2	0.020
15	128	N,N-Dimethyl-5-cyclohexylpentylamine	0.26	0.03	3	195.33	27.82	3	0.001
	129	N-Benzyl-N'-(5-phenylpentyl)-piperazine	0.41	0.04	2	11.32	1.70	3	.036
	134	N-Benzyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine	127.90	25.04	5	470.75	37.40	4	0.272
	137	N-Benzyl-N-methyl-5-phenylpentylamine	0.19	0.04	2	12.55	0.15	2	0.015
	140	N-(3-phenylpropyl)piperidine	48.94	4.39	3	146.50	3.50	2	

-130-

TABLE 11

COMPOUND	SIGMA-1 (n M) Ki vs. 3H-PENTAZOCINE			SIGMA-2 (n M) Ki vs. 3H-DTG +PENTAZOCINE			S1/S2 Ratio
	MEAN	SEM	n	MEAN	SEM	n	
(+)-3-PPP	47.67	1.45	3	493.00	40.07	3	0.097
(-)-3-PPP	312.33	17.89	3	1261.00	87.32	3	0.248
(+)-Butaclamol	583.67	177.34	3	1399.67	86.15	3	0.417
(-)-Butaclamol	69.33	14.89	3	3422.67	723.43	3	0.020
Chlorpromazine	336.00	23.58	4	714.00	201.00	2	0.471
Dextromethorphan	522.00	239.08	3	14990.00	2653.32	3	0.035
Fluoxetine	56.75	8.8353	4	10000	0	3	
Fluphenazine	31.67	7.62	3	67.33	8.50	4	0.470
Ifenprodol	5.83	0.64	3	2.39	0.39	3	2.444
(+)-Pentazocine	1.60	0.09	4	906.33	27.39	3	0.002
(-)-Pentazocine	114.00	20.30	3	242.33	8.97	3	0.470
Perphenazine	25.10	7.45	3	80.35	0.05	2	0.312
Pramiracetam	10000.00	0.00	3	10000.00	--	1	1.000
Remoxipride	312.00	36.58	4	1617.00	267.38	3	0.193
Risozazole	1156.00	183.06	3	264.67	86.11	3	4.368
(+)-SKF 10047	148.00	19.09	3	16173.25	3447.26	4	0.009
(-)-SKF 10047	3640.00	825.19	3	9131.25	1574.11	4	0.399
Thioridazine	520.50	81.22	4	334.50	82.50	2	1.556
Trifluoperazine	561.25	135.80	4	160.50	47.50	2	3.497
Spiperone	10000.00	--	1	--	--	0	--

EXAMPLE 8 Structure-Activity Analysis with Regard to Sigma-1

As shown in Table 12, compound 1 binds with an IC_{50} value of 10.8 nM at sigma-1. Substitution on the phenyl A ring has essentially no effect on binding.




Table 12			
Cmp. #	X	Isomer	IC_{50} (nM)
1	H	R(-)	10.8
43	H	S(+)	--
20	3-Br	(±)	9.6
21	4-Br	(±)	12.0
22	4-I	R(-)	17.7

23	3-CF ₃	(±)	8.8
83	4-OEt	(±)	3.4


Replacement of the phenyl-A with 1-naphthyl has no effect on binding. (Compare compound no. 1 to compound no. 16 (Table 13)). However, replacement of the 1-naphthyl group with a 2-naphthyl group reduces sigma-1 affinity about three fold. (Compare compound no. 16 to compound no. 17 (Table 13)). In addition, replacement of phenyl-B with 1-naphthyl or 2-naphthyl has little effect on binding. (Compare compound no. 1 to compound nos. 11 and 12 (Table 13)).

Table 13

Cmp.	Structure	Sigma-1 IC ₅₀ (nM)
------	-----------	-------------------------------

15


1



10.8

20


16



9.3

25


17



31.0

30


11



8.6

35

12



5.7

Increasing the length of the N-phenyl-B alkyl chain of compound No. 1 from 3 to 4 methylene groups doubles affinity to the sigma-1 receptor. A further increase to 5 methylene groups substantially increases affinity. (Compare compound no. 1 to compound nos. 10 and 45 (Table 14)). Removal of the σ -methyl group has little effect on sigma-1 binding. (Compare compound no. 1 to compound no. 87 (Table 14)).




Table 14

Cmp. #	n	Isomer	IC ₅₀ (nM)
1 R = CH ₃	3	R(-)	10.8
10 R = CH ₃	4	(±)	4.7
44 R = CH ₃	4	S(+)	--
45 R = CH ₃	5	S(+)	.88
87 R = H	3		11.3

Holding the methylene groups linking the phenyl-B ring constant at four and removing the σ -methyl group does not affect substantially the affinity to sigma-1. (Compare cmp. nos. 71 and 10 (N-phenyl-A; n=2), which have binding affinities of 2.6 and 4.7 nM, respectively). Shortening the number of methylene groups linking the phenyl-A substituent to one decreases affinity. (Compare cmp. nos. 71 and 75, which have binding affinities of 2.6 and 9.6, respectively.) In contrast, increasing the number of methylene groups linking the phenyl-A ring to five increases affinity. (Compare cmp. nos. 71 and 76, which have binding affinities of 2.6 and 0.48 nM, respectively.)

Since five methylenes between the phenyl-A ring and the amine is associated with high binding to sigma-1, a homologous series of related analogs were examined (Table 15).




Table 15			
cmp. #	n	R	IC ₅₀ (nM)
76	4	H	0.48
111	3	H	0.28
89	2	H	0.17
77	1	H	0.32
137	1	CH ₃	0.19

As is evident from Table 15, the length of the N-phenyl-B chain has little effect on binding to the sigma-1 site. In addition, elimination of the N-phenyl-B ring has little effect on sigma-1 receptor binding. (Cmp. #112, N-methyl-N-propyl-5-phenylpentylamine, IC₅₀=0.29 nM). However, an increase in the phenyl-A chain length results in a decrease in affinity to sigma-1 (compare cmp. nos. 90, N-benzyl-7-phenylheptylamine, and cmp. 91, N-phenethyl-7-phenylheptylamine which have binding affinities at sigma-1 of 2.3 and 1.5 nM, respectively).


Reduction of the phenyl-A ring of cmp. no. 77 (N-benzyl-5-phenylpentylamine, IC₅₀=0.32) has little effect on sigma-1 binding affinity (cmp. #94, N-benzyl-5-cyclohexylpentylamine, IC₅₀=0.81).

Removal of the aromatic phenyl-B ring of the A-ring cyclohexyl compounds is also well tolerated. However, the tertiary amines appear to have enhanced affinity compared to the primary and secondary amines (Table 16).

Table 16



Cmp. #	Structure	Sigma-1 IC ₅₀ (nM)
--------	-----------	-------------------------------

94



0.81

-134-



15 The most selective sigma-1 binding compound is N,N-dimethyl-5-cyclohexylpentylamine (cmp. # 128, sigma-2/sigma-1 selectivity of 650). Other sigma-1 selective ligands include N-methyl-5-cyclohexylpentylamine (cmp. # 127, sigma-2/sigma-1 selectivity of 51), 5-cyclohexylpentylamine (cmp. # 126, sigma-2/sigma-1 selectivity of 13), N-methyl-N-propyl-5-phenylpentylamine (cmp. # 112, sigma-2/sigma-1 selectivity of 137), N-benzyl-7-phenylheptylamine (cmp. # 90, sigma-2/sigma-1 selectivity of 126), N-(5-phenyl)pentylpiperidine (cmp. # 124, sigma-2/sigma-1 selectivity of 100), and N,N-dimethyl-5-phenylpentylamine (cmp. # 125, sigma-2/sigma-1 selectivity of 96).

EXAMPLE 9 Structure-Activity Analysis with Regard to Sigma-2

30 Many alterations in structure do not improve the binding of the compounds of the present invention to the sigma-2 site. Compound #1 binds at the sigma-2 site with modest affinity (60 nM). Aromatic substitution in the phenyl-A ring has little effect on binding at sigma-2 (Table 17).




Table 17			
cmp. #	X	Isomer	IC ₅₀ (nM)
1	H	R(-)	60
43	H	S(+)	--
20	3-Br	(±)	26
21	4-Br	(±)	39
22	4-I	R(-)	76
23	3-CF ₃	(±)	21
83	4-OEt	(±)	34

However, replacement of phenyl-A with a naphthyl ring decreases affinity to sigma-2 several fold. (Compare cmp. #1, IC₅₀ = 60 with cmp. # 16, N-(3-phenylpropyl)-1-(1-naphthyl)-2-isopropylamine, and #17, N-(3-phenylpropyl)-1-(2-naphthyl)-2-isopropylamine, which have binding affinities of 150 and 220 nM, respectively.) In addition, replacement of phenyl-B with a naphthyl ring reduces affinity about 4 fold. (Compare cmp. #1, IC₅₀ = 60 with cmp. # 11, N-(3-(1-naphthyl)propyl)-1-phenyl-2-isopropylamine, and #1, N-(3-(2-naphthyl)propyl)-1-phenyl-2-isopropylamine, which have binding affinities of 280 and 260 nM, respectively.) Removal of the α -methyl group of cmp. # 1 has essentially no effect (IC₅₀ of cmp. no. 87, N-(3-phenylpropyl)-2-phenethylamine, = 90 nM). Increasing the phenyl-B chain length from 3 to 4 also has little effect (IC₅₀ of cmp. 71, N-(4-phenylbutyl)-2-phenethylamine = 120 nM). Decreasing the phenyl-A alkyl chain length by one methylene slightly decreases affinity (IC₅₀ of cmp. # 75, N-(4-phenylbutyl)benzylamine, = 160 nM).

However, it is possible to improve binding affinity to sigma-2 by keeping the phenyl-B chain at 4 methylenes and increasing the phenyl-A chain to 5 methylenes to give a 3-fold increase in affinity (Table 18). An alkyl chain of n=3 appears to be optimal. In addition, N-methylation triples affinity.




Table 18			
cmp. #	n	R	IC ₅₀ (nM)
76	4	H	48
111	3	H	9.8
89	2	H	15
77	1	H	34
37	1	CH ₃	12

Removal of the phenyl-B ring results in a reduction in sigma-2 affinity. (Compare cmp. 111, N-(3-phenylpropyl)-5-phenylpentylamine, with N-methyl-N-propyl-5-phenylpentylamine, IC₅₀=40 nM).


The most selective sigma-2 ligand of the present invention is N-phenyl-N'-(3-(1-phthalimido)propyl)piperazine (cmp. #96) which has a sigma-1/sigma-2 ratio of 87. Other sigma-2 selective ligands include N-(4-phthalimido)butyl-N'-(o-methoxyphenyl)piperazine (cmp. #120; sigma-1/sigma-2 ratio of 7) and N-(4-phthalimido)butyl-N'-phenylpiperazine (cmp. #97; sigma-1/sigma-2 ratio of 10).

Having now fully described this invention, it will be understood by those of skill in the art that the same can be practiced within a wide range of equivalent conditions, formulations, structural variations and other parameters without affecting the scope of the invention or any embodiment thereof.

-137-

WHAT IS CLAIMED IS:

1. A method of treating a human being suffering from drug abuse, gastrointestinal disorder or depression, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl;

R is hydrogen or C₁-C₆ alkyl;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo and =O; or

R and R¹ together form a morpholino, piperazinyll or piperidinyl ring;



n is 0-5;

W is -(CH₂)_p- or -H H-, wherein p is 1-3;

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-,

-138-



5 $-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S; or

C_1-C_6 alkyl (wherein Z is hydrogen);


Z is hydrogen, cycloalkyl, aryl, an aryl-substituted carboxylic acid group, or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, 10 C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkylamino, C_2-C_{15} dialkylamino, hydroxy, carbamoyl, C_1-C_6 N -alkylcarbamoyl, C_2-C_{15} N,N -dialkylcarbamoyl, nitro, C_2-C_{15} dialkylsulfamoyl or an ortho methylene-dioxy group; 20

wherein said compound exhibits high binding activity with respect to the sigma receptor.

2. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound of the formula: 25

30

35



wherein:

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl;

R is hydrogen or C₁-C₆ alkyl;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo and =O; or R and R¹ together form a morpholino, piperaziny or piperidiny ring;

n is 0-5;

W is -(CH₂)_p- or -H H-, wherein p is 1-3;

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-;

$$\begin{array}{c} \text{O} \\ \parallel \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-;} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen);


Z is cycloalkyl or an aryl-substituted carboxylic acid group, wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy,

-140-

amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or an ortho methylene-dioxy group;

wherein said compound exhibits high binding activity with respect to the sigma receptor.

3. A method of treating a human being suffering from drug abuse, gastrointestinal disorder or depression, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

R is hydrogen or C₁-C₆ alkyl;

T is aryl, heteroaryl, substituted aryl, substituted heteroaryl, or cycloalkyl, wherein said substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl, and an ortho methylenedioxy group;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, bromo, chloro, iodo and =O; or

R and R¹ together form a morpholino, piperazinyll or piperidinyl ring;


n is 0-5;

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein r is 0-3;

-(CH₂)_r-CH=CH-(CH₂)_r-;

-141-



$$-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-, \text{ wherein each } r \text{ is independently } 0-3 \text{ and}$$

5 wherein Y is O or S; or

$\text{C}_1\text{-C}_6$ alkyl (wherein Z is hydrogen);

Z is hydrogen, cycloalkyl, aryl, an aryl-substituted carboxylic acid group, or aryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_2\text{-C}_6$ dialkoxymethyl, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_3\text{-C}_{15}$ dialkylaminoalkyl, carboxy, carboxamido, 10 $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ haloalkylthio, allyl, aralkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, aroyl, aralkoxy, $\text{C}_2\text{-C}_6$ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, $\text{C}_3\text{-C}_6$ heterocycloalkyl, $\text{C}_1\text{-C}_6$ alkylthio, $\text{C}_1\text{-C}_6$ alkylsulfonyl, $\text{C}_1\text{-C}_6$ haloalkylsulfonyl, 15 $\text{C}_1\text{-C}_6$ alkylsulfinyl, $\text{C}_1\text{-C}_6$ haloalkylsulfinyl, arylthio, $\text{C}_1\text{-C}_6$ haloalkoxy, amino, $\text{C}_1\text{-C}_6$ alkylamino, $\text{C}_2\text{-C}_{15}$ dialkylamino, hydroxy, carbamoyl, $\text{C}_1\text{-C}_6$ N-alkylcarbamoyl, $\text{C}_2\text{-C}_{15}$ N,N-dialkylcarbamoyl, nitro, $\text{C}_2\text{-C}_{15}$ dialkylsulfamoyl, or an ortho methylene-dioxy group;


20 wherein said compound exhibits high binding activity with respect to the sigma receptor.

4. The method of claim 3, wherein said compound is selected from the group consisting of N-phenethyl-1-phenylisopropylamine, N-(3-phenylpropyl)-1-phenyl-isopropylamine, 25 N-(2-phenoxy)ethyl)-1-phenylisopropylamine, N-(3-phenyl-3-propanon-1-yl)-1-phenylisopropylamine, N-(4-phenylbutyl)-1-phenylisopropylamine, N-(3-(1-naphthyl)propyl)-1-phenyl-isopropylamine, N-(3-(2-naphthyl)propyl)-1-phenylisopropylamine, 30 N-methyl-N-(3-(2-naphthyl)propyl)-1-phenylisopropylamine, N-(3-phenyl-2-propyn-1-yl)-1-phenyl-isopropylamine, N-(3-phenylpropyl)-1-(4-hydroxyphenyl)isopropylamine, N-(3-phenylpropyl)-1-(4-methoxyphenyl)isopropylamine, N-(3-phenylpropyl)-1-(3-bromophenyl)isopropylamine, N-(3-phenylpropyl)-1-(4-bromophenyl)isopropylamine, N-(3-phenylpropyl)-1-(3,4-dichlorophenyl)isopropylamine, N-(3-phenylpropyl)-1-(4-iodophenyl)-isopropyl)-amine, N-(3-phenylpropyl)-1-(3-trifluoromethyl-

phenyl)isopropylamine, N-(2-phenethyl)-N-methyl-1-phenyl-
isopropylamine, N-(3-phenylpropyl)-1-phenylpropan-1-one-2-amine,
N-(2-indanyl)-3-phenylpropylamine, N,N-di-(3-phenylpropyl)amine,
N-(2-(1-naphthyl)ethyl)-1-phenylisopropylamine, N-(2-(2-
5 naphthyl)ethyl)-1-phenylisopropylamine, N-(2-(1-naphthyl)propyl)-
1-phenylisopropylamine, N-(2-(2-naphthyl)propyl)-1-phenyliso-
propylamine, N-(4-phenylbutyl)-1-(2,5-dimethoxyphenyl)iso-
propylamine, N-(4-phenylbutyl)-1-(4-bromo-2,5-dimethoxyphenyl)-
isopropylamine, N-methyl-N-(4-phenylbutyl)-1-phenylisopropyl-
10 amine, N-methyl-N-(5-phenylpentyl)-1-phenylisopropylamine, N-(3-
phenylpropyl)-1-(4-ethoxyphenyl)isopropyl-amine, N-(3-phenyl-
propyl)-1-(4-propylphenyl)isopropylamine, N-(3-phenylpropyl)-1-
(4-benzoyloxyphenyl)isopropylamine, N-methyl-N-(3-phenylpropyl)-1-
(4-propylphenyl)isopropylamine, N-(3-phenylpropyl)-1-phenyl-2-
15 pentylamine, N-(4-phenylbutyl)-1-phenyl-2-pentylamine, N,N-di-(2-
ethylphenyl)methylamine, N,N-dibenzylamine, N-(3-phenylpropyl)-N-
(6-phenylhexyl)amine, N-(3-phenylpropyl)-N-(5-phenylpentyl)amine,
N-propyl-N-methyl-5-phenylpentylamine, N-methyl-N-(3-phenyl-
propyl)-1-phenylisopropylamine, N-methyl-N-(3-methyl-2-butenyl)-
20 1-phenyliso-propylamine, N-methyl-N-(3-methylbutyl)-1-phenyl-
isopropylamine, N-methyl-N-(3-phenylpropyl)-1-phenyl-2-
pentylamine, N-methyl-N-(3-methylbutyl)-1-isopropylamine, N-
methyl-N-(3-phenylbutyl)-1-phenyl-2-pentylamine, N-propyl-N-(3-
phenyl)propyl)-1-phenyl-2-propylamine, N-benzyl-N-(3-phenyl)-
25 propyl)-1-phenyl-2-propylamine, N-phenyl-(5-phenyl)pentylamine,
N-methyl-N-(3-phenyl)propyl-5-phenylpentylamine, N-(2-(o-methyl-
phenyl)ethyl)-5-phenylpentylamine, N-(2-(m-methylphenyl)ethyl)-5-
phenylpentylamine, N-(2-(p-methylphenyl)ethyl)-5-phenylpentyl-
amine, N-benzyl-5-phenylpentylamine, N-benzyl-N-methyl-5-phenyl-
30 pentylamine, N-(2-(3-hydroxyphenyl)ethyl)-5-phenylpentylamine, N-
(2-(2-hydroxyphenyl)ethyl)-5-phenylpentylamine, N,N'-diethyl-2-
(diphenylacetoxy)ethylamine, N,N'-diethyl-2-(9-fluorene-carboxy)-
ethylamine, N,N-Dimethyl-5-phenylpentylamine, N-Benzyl-N-(3-
phenylpropyl)-1-phenyl-2-propylamine, N-Benzyl-N-methyl-5-
35 phenylpentylamine, N-Benzyl-5-phenylpentylamine, and N-(2-
phenethyl)-N-methylpentylamine.

-143-

5. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

R is hydrogen or C₁-C₆ alkyl;

T is aryl, heteroaryl, substituted aryl, substituted heteroaryl, or cycloalkyl, wherein said substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl, and an ortho methylenedioxy group;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, bromo, chloro, iodo and =O; or R and R¹ together form a morpholino, piperazinyll or piperidinyl ring;

n is 0-5;

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein r is 0-3;

-(CH₂)_r-CH=CH-(CH₂)_r-;

$\begin{array}{c} \text{O} \\ || \\ -(\text{CH}_2)_r-\text{C}-(\text{CH}_2)_r-; \end{array}$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen);

-144-

5 Z is cycloalkyl or an aryl-substituted carboxylic acid group wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl, or an ortho methylene-dioxy group;

10 wherein said compound exhibits high binding activity with respect to the sigma receptor.

15


6. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound selected from the group consisting of N-(3-phenyl-propyl)-1-(4-propylphenyl)isopropylamine, N-(3-phenylpropyl)-1-(4-benzoxyphe-
20 nyl)isopropylamine, N-methyl-N-(3-phenylpropyl)-1-(4-propylphenyl)isopropylamine, N-(3-phenylpropyl)-1-phenyl-2-pentylamine, N-(4-phenylbutyl)-1-phenyl-2-pentylamine, N,N-di-(2-ethylphenyl)methylamine, N,N-dibenzylamine, N-(3-phenylpropyl)-N-(6-phenylhexyl)amine, N-(3-phenylpropyl)-N-(5-phenylpentyl)amine, N-propyl-N-methyl-5-phenylpentylamine, N-methyl-N-(3-phenylpropyl)-1-phenylisopropylamine, N-methyl-N-(3-methyl-2-butenyl)-1-phenyliso-propylamine, N-methyl-N-(3-methylbutyl)-1-phenylisopropylamine, N-methyl-N-(3-phenylpropyl)-1-phenyl-2-pentylamine, N-methyl-N-(3-methylbutyl)-1-isopropylamine, N-methyl-N-(3-phenylbutyl)-1-phenyl-2-pentylamine, N-propyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine, N-benzyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine, N-phenyl-(5-phenyl)pentylamine, N-methyl-N-(3-phenyl)propyl-5-phenylpentylamine, N-(2-(o-methylphenyl)ethyl)-5-phenylpentylamine, N-(2-(m-methylphenyl)ethyl)-5-phenylpentylamine, N-(2-(p-methylphenyl)ethyl)-5-phenylpentyl-

25
30
35

-145-

amine, N-benzyl-5-phenylpentylamine, N-benzyl-N-methyl-5-phenylpentylamine, N-(2-(3-hydroxyphenyl)ethyl)-5-phenylpentylamine, N-(2-(2-hydroxyphenyl)ethyl)-5-phenylpentylamine, N,N'-diethyl-2-(diphenylacetoxy)ethylamine, N,N'-diethyl-2-(9-fluorene-carboxy)-ethylamine, N,N-Dimethyl-5-phenylpentylamine, N-Benzyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine, N-Benzyl-N-methyl-5-phenylpentylamine, N-Benzyl-5-phenylpentylamine, N-(2-phenethyl)-N-methylpentylamine, N-(5-phenylpentyl)-4-benzylpiperidine and N-(5-phenylpentyl)-4-benzyl-4-hydroxy-piperidine.

7. A method of treating a human being suffering from drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

R is hydrogen or C₁-C₆ alkyl;

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅

-146-

dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O; or

5 R and R¹ together form a morpholino, piperazinyll or piperidinyl ring;

n is 1-3;

p is 1-3;

X is -(CH₂)_q-, wherein q is 1-6,

10 -(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-,

O
||
-(CH₂)_r-C-(CH₂)_r-,

15 -(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and wherein Y is O or S; or

C₁-C₆ alkyl;

Z is hydrogen, cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

30 wherein said compound exhibits a high binding activity with respect to the sigma receptor.


8. The method of claim 7, wherein said compound is selected from the group consisting of N-(2-indanyl)-1-phenylisopropylamine, N-(2-indanyl)-3-phenylpropylamine, N-(4,5-benzocycloheptyl)-1-phenylisopropylamine, N-(3,4-benzocycloheptyl)-3-(phenyl)propylamine, N-(4,5-benzocycloheptyl)-3-

35

-147-

phenylpropylamine, N-(3,4-benzocyclohexyl)-1-phenylisopropylamine and N-3,4-benzocyclohexyl-3-phenylpropylamine.

9. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

R is hydrogen or C₁-C₆ alkyl;

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O; or



R and R¹ together form a morpholino, piperazinyll or piperidinyl ring;

n is 1-3;


p is 1-3;

X is -(CH₂)_q-, wherein q is 1-6,

-148-



5



$-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S; or

$\text{C}_1\text{-C}_6$ alkyl;

10

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_2\text{-C}_6$ dialkoxymethyl, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_3\text{-C}_{15}$ dialkylaminoalkyl, carboxy, carboxamido, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ haloalkylthio, allyl, aralkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, aroyl, aralkoxy, $\text{C}_2\text{-C}_6$ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, $\text{C}_3\text{-C}_6$ heterocycloalkyl, $\text{C}_1\text{-C}_6$ alkylthio, $\text{C}_1\text{-C}_6$ alkylsulfonyl, $\text{C}_1\text{-C}_6$ haloalkylsulfonyl, $\text{C}_1\text{-C}_6$ alkylsulfinyl, $\text{C}_1\text{-C}_6$ haloalkylsulfinyl, arylthio, $\text{C}_1\text{-C}_6$ haloalkoxy, amino, $\text{C}_1\text{-C}_6$ alkylamino, $\text{C}_2\text{-C}_{15}$ dialkylamino, hydroxy, carbamoyl, $\text{C}_1\text{-C}_6$ N -alkylcarbamoyl, $\text{C}_2\text{-C}_{15}$ N,N -dialkylcarbamoyl, nitro, $\text{C}_2\text{-C}_{15}$ dialkylsulfamoyl or methylenedioxy;


20

wherein said compound exhibits a high binding activity with respect to the sigma receptor.

25

10. A method of treating a human being suffering from drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a compound of the formula:

30



35

wherein:

R^2 is selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_2\text{-C}_6$ dialkoxymethyl, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_3\text{-C}_{15}$ dialkylaminoalkyl, carboxy,

-149-

carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

X is -(CH₂)_q-, wherein q is 1-6,

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-,

$$\begin{array}{c} \text{O} \\ || \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and

wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen);


Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

V is N or CM, wherein M is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, hydroxy, fluoro, chloro, bromo or represents one half of a double bond with a neighboring endocyclic carbon;

wherein said compound exhibits a high binding activity with respect to the sigma receptor.

11. The method of claim 10, wherein V is N, said compound having the formula

-150-




5

12. The method of claim 11, wherein said compound is selected from the group consisting of N-(3-trifluoromethylphenyl)-N'-phenethylpiperazine, N-(3-trifluoromethylphenyl)-N'-(4-phenylbutyl)piperazine, N-(3-chlorophenyl)-N'-benzylpiperazine, N-(3-chlorophenyl)-N'-(3-phenylpropyl)piperazine, N-(3-chlorophenyl)-N'-phenethylpiperazine, N-phenyl-N'-(3-phenylpropyl)piperazine, N-phenyl-N'-(3-phenylbutyl)piperazine, N-(2-naphthyl)-N'-(3-phenylpropyl)-piperazine, N-phenyl-N'-(3-(2-naphthyl)propyl)-piperazine, N-phenyl-N'-propylpiperazine, N-(4-chlorophenyl)-N'-(3-phenylpropyl)piperazine, N-benzyl-N'-(4-phenylbutyl)-piperazine, N-phenyl-N'-(4-phthalimidobutyl)-piperazine, and N-phenyl-N'-(5-phthalimidopentyl)piperazine.

15

20

13. The method of claim 11, wherein V is CM, said compound having the formula



25

30

14. The method of claim 13, wherein said compound is selected from the group consisting of N-phenethyl-4-phenylpiperidine, N-phenethyl-4-phenyl-4-hydroxypiperidine, N-(3-phenylpropyl)-4-phenylpiperidine, N-(3-phenylpropyl)-4-phenyl-4-hydroxypiperidine, N-(5-phenylpentyl)-4-phenylpiperidine, N-(4-phenylbutyl)-4-phenylpiperidine, N-(3-phenylpropyl)-4-(4-chlorophenyl)-1,2,5,6-tetrahydropyridine, N-(4-phenylbutyl)-4-(4-chlorophenyl)-1,2,5,6-tetrahydropyridine, N-(5-phenylpentyl)-4-


35

40

-151-

benzylpiperidine and N-(5-phenylpentyl)-4-benzyl-4-hydroxy-piperidine.

15. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

R^2 is selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxy-methyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkylamino, C_2-C_{15} dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro and C_2-C_{15} dialkylsulfamoyl;

X is $-(CH_2)_q-$, wherein q is 1-6,

$-(CH_2)_r-C\equiv C-(CH_2)_r-$,

$-(CH_2)_r-CH=CH-(CH_2)_r-$,

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$,

$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and

wherein Y is O or S; or

C_1-C_6 alkyl (wherein Z is hydrogen);

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-


-152-

C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;


V is N or CM, wherein M is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, hydroxy, fluoro, chloro, bromo or represents one half of a double bond with a neighboring endocyclic carbon;

wherein said compound exhibits a high binding activity with respect to the sigma receptor.

16. The method of claim 15, wherein V is N, said compound having the formula




17. The method of claim 15, wherein V is CM, said compound having the formula



18. A method of treating a human being suffering from drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a compound of the formula:

-153-



wherein:

R^3 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_2 - C_6 dialkoxymethyl, C_3 - C_{15} dialkylaminoalkyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, C_2 - C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl;

X is $-(CH_2)_q-$, wherein q is 1-6,

$-(CH_2)_r-C\equiv C-(CH_2)_r-$,

$-(CH_2)_r-CH=CH-(CH_2)_r-$,

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$,

$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S; or

C_1 - C_6 alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1 - C_6 alkoxy, C_2 - C_6 dialkoxymethyl, C_1 - C_6 alkyl, cyano, C_3 - C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkylthio, allyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, aralkoxy, C_2 - C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl, C_1 - C_6 alkylthio, C_1 - C_6 alkylsulfonyl, C_1 - C_6 haloalkylsulfonyl, C_1 - C_6 alkylsulfinyl, C_1 - C_6 haloalkylsulfinyl, arylthio, C_1 - C_6 haloalkoxy, amino, C_1 - C_6 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C_1 - C_6 N-alkylcarbamoyl, C_2 - C_{15} N,N-dialkylcarbamoyl, nitro, C_2 - C_{15} dialkylsulfamoyl or methylenedioxy;


V is N or CM, wherein M is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, hydroxy, fluoro, chloro, bromo or represents one half of a double bond with a neighboring endocyclic carbon;

wherein said compound exhibits a high binding activity with respect to the sigma receptor.

-154-

19. The method of claim 18, wherein said compound is selected from the group consisting of N-methyl-N'-(4-phenyl-3-(E)butenyl)piperazine, N-methyl-N'-(4-phenyl-3-(Z)butenyl)piperazine, N-methyl-N'-(4-(3-trifluoromethylphenyl)-3-(Z)butenyl)piperazine, N-methyl-N'-(4-phenylbutyl)piperazine, N-benzyl-N'-(4-phthalimidobutyl)piperazine, N-(2-methoxyphenyl)-N'-(4-phthalimidobutyl)piperazine, N-(5-phenylpentyl)-4-benzylpiperidine, and N-(5-phenylpentyl)-4-benzyl-4-hydroxy-piperidine.

20. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

R^3 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_2 - C_6 dialkoxymethyl, C_3 - C_{15} dialkylaminoalkyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, C_2 - C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl;

X is $-(CH_2)_q-$, wherein q is 1-6,

$-(CH_2)_r-C\equiv C-(CH_2)_r-$,

$-(CH_2)_r-CH=CH-(CH_2)_r-$,

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$,

$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and

wherein Y is O or S; or

C_1 - C_6 alkyl (wherein Z is hydrogen);

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1 - C_6 alkoxy, C_2 - C_6 dialkoxymethyl, C_1 - C_6 alkyl, cyano, C_3 - C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkylthio, allyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, aralkoxy, C_2 - C_6 carboxylic acyl, aryl, substituted


-155-

aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

V is N or CM, wherein M is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, hydroxy, fluoro, chloro, bromo or represents one half of a double bond with a neighboring endocyclic carbon;

wherein said compound exhibits a high binding activity with respect to the sigma receptor.

21. A method of treating a human being suffering from a central nervous system disorder, drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein:

R⁴ is hydrogen or an aryl group substituted with a group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₂-C₆ dialkoxymethyl, C₃-C₁₅ dialkylaminoalkyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl;

R⁵ is hydrogen or hydroxy;

X is -(CH₂)_q-, wherein q is 1-6,

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-,

$$\begin{array}{c} \text{O} \\ \parallel \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and

wherein Y is O or S; or

-156-

C₁-C₆ alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;


wherein said compound exhibits a high binding activity with respect to the sigma receptors.

22. The method of claim 21, wherein said compound is selected from the group consisting of N-(5-phenylpentyl)-piperidine, N-(8-phenylheptyl)piperidine, N-(5-(4-methoxyphenyl)pentyl)piperidine, N-(3-phenylpropyl)piperidine, N-(5-cyclohexyl)pentylpiperidine, N-benzylpiperidine, N-(2-phenethyl)-4-hydroxy-4-phenylpiperidine, N-(2-phenethyl)-4-hydroxy-4-t-butylpiperidine, N-(5-(4-chlorophenyl)-5-pentanone-1-yl)piperidine, N-(5-(4-chlorophenyl)-5-pentanone-1-yl)-4-phenylpiperidine, N-(5-(4-methoxyphenyl)-5-pentanone-1-yl)piperidine, N-(5-(4-methoxyphenyl)-5-pentanone-1-yl)-4-phenylpiperidine, N-(5-(4-methoxyphenyl)pentyl)-4-phenylpiperidine, N-(5-phenyl-5-pentanone-1-yl)-4-phenylpiperidine, N-(5-(4-chlorophenyl)-pentyl)-4-phenylpiperidine, N-(5-(3-methoxyphenyl)-5-pentanone-1-yl)piperidine, N-(5-(3-chlorophenyl)-5-pentanone-1-yl)piperidine, N-(5-(3-chlorophenyl)-5-pentanone-1-yl)-4-phenylpiperidine, N-(5-(3-methoxyphenyl)-5-pentanone-1-yl)-4-phenylpiperidine, N-(4-(4-fluorophenyl)-4-butanone-1-yl)piperidine, N-(5-(4-fluorophenyl)-5-pentanone-1-yl)piperidine, N-(5-(4-fluorophenyl)-5-pentanone-1-yl)-4-phenylpiperidine, N-(5-(4-fluorophenyl)-5-pentanone-1-yl)-4-(3-chlorophenyl)-4-hydroxypiperidine, N-(5-(4-chlorophenyl)-5-

-157-

pentanon-1-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)-piperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)-piperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(chlorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(chlorophenyl)piperidine, N-(5-(3,4-dichlorophenyl)-5-pentanon-1-yl)-4-(chlorophenyl)-piperidine, N-(5-cyclopentylpentan-5-on-1-yl)piperidine and N-(5-(3,4-methylenedioxyphenyl)penta-2,4-dienyl)piperidine.

23. A method of treating a human being suffering from a central nervous system disorder, drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



wherein,

X is $-(CH_2)_q-$, wherein q is 1-6,

$-(CH_2)_r-C\equiv C-(CH_2)_r-$,

$-(CH_2)_r-CH=CH-(CH_2)_r-$,

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$,

$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and

wherein Y is O or S; or

C_1-C_6 alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio,


-158-

allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

24. The method of claim 23, wherein said compound is N-(5-phenyl)pentyl-3-azabicyclo[3.2.2]nonane.

25. A method of treating a human being suffering from a central nervous system disorder, drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a tropane derivative having the formula:



wherein,

R⁴ is hydrogen or an aryl group substituted with a group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkenyl, C₂-C₆ dialkoxymethyl, C₃-C₁₅ dialkylaminoalkyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl;


R⁵ is hydrogen or hydroxy;

X is -(CH₂)_q-, wherein q is 1-6,

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-,

-159-



5 $-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S; or


$\text{C}_1\text{-C}_6$ alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_2\text{-C}_6$ dialkoxymethyl, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_3\text{-C}_{15}$ dialkylaminoalkyl, carboxy, carboxamido, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ haloalkylthio, allyl, aralkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, aroyl, aralkoxy, $\text{C}_2\text{-C}_6$ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, $\text{C}_3\text{-C}_6$ heterocycloalkyl, $\text{C}_1\text{-C}_6$ alkylthio, $\text{C}_1\text{-C}_6$ alkylsulfonyl, $\text{C}_1\text{-C}_6$ haloalkylsulfonyl, $\text{C}_1\text{-C}_6$ alkylsulfinyl, $\text{C}_1\text{-C}_6$ haloalkylsulfinyl, arylthio, $\text{C}_1\text{-C}_6$ haloalkoxy, amino, $\text{C}_1\text{-C}_6$ alkyl-amino, dialkylamino, hydroxy, carbamoyl, $\text{C}_1\text{-C}_6$ N-alkylcarbamoyl, $\text{C}_2\text{-C}_{15}$ N,N-dialkylcarbamoyl, nitro, $\text{C}_2\text{-C}_{15}$ dialkylsulfamoyl or methylenedioxy;

20 wherein said compound exhibits a high binding activity with respect to the sigma receptors.

26. The method of claim 25, wherein said compound is N-(5-phenyl)pentyl-4-phenyltropan-4-ol.

25 27. A method of treating a human being suffering from drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a compound of the formula:



wherein

a is 1-8;

b is 1-8;

-160-

R is hydrogen or C₁-C₆ alkyl;

R² is independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;


wherein said compound exhibits high binding activity with respect to the sigma receptor.

28. The method of claim 27 wherein said compound is selected from the group consisting of N-(4-phenylbutyl)phenethylamine, N-(4-phenylbutyl)-3-phenylpropylamine, N-(4-phenylbutyl)-4-phenylbutylamine, N-(4-phenylbutyl)benzylamine, N-(4-phenylbutyl)-5-phenylpentylamine, N-(5-phenylpentyl)benzylamine, N-(3-phenylpropyl)phenethylamine, N-(5-phenylpentyl)phenethylamine, N-(7-phenylheptyl)benzylamine, N-(7-phenylheptyl)-phenethylamine, N-methyl-N-(2-phenethyl)-1-phenylisopropylamine, N-methyl-N-(5-phenylpentyl)-1-phenylisopropylamine, N-methyl-N-(3-phenylpropyl)-1-(4-propylphenyl)isopropylamine, N-phenyl-(5-phenyl)pentylamine, N-methyl-N-(3-phenylpropyl)-5-phenylpentylamine, N-benzyl-N-methyl-5-phenylpentylamine, N-(2-(o-methoxyphenyl)ethyl)-5-phenylpentylamine, N-(2-(m-methoxyphenyl)ethyl)-5-phenylpentylamine, N-(2-(p-methoxyphenyl)ethyl)-5-phenylpentylamine, N-benzyl-5-phenylpentylamine, N-(2-(m-hydroxyphenyl)ethyl)-5-phenylpentylamine, and N-(2-(o-hydroxyphenyl)ethyl)-5-phenylpentylamine.

29. A method of treating a human being suffering from drug abuse, gastrointestinal disorder, or depression, which

-161-

comprises administering to said human a therapeutically effective amount of a compound having the Formula:



wherein

10 a is 1-8;

b is 1-8;

R is hydrogen or C₁-C₆ alkyl;

15 R² is independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkyl-

20 sulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl; and

25 wherein said compound exhibits high binding activity with respect to the sigma receptor.

30 30. The method of claim 29, wherein said compound is N-(3-phenylpropyl)-2-(2-naphthyl)ethylamine.

31. A method of treating a human being suffering from drug abuse, gastrointestinal disorder or depression, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



wherein;

5 Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, 10 allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

X is -(CH₂)_q-, wherein q is 1-6,

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-,

20

$$\begin{array}{c} \text{O} \\ || \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and

wherein Y is O or S; or

25


C₁-C₆ alkyl;

Z is hydrogen, cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

-163-

wherein said compound exhibits high binding activity with respect to the sigma receptor.

32. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



wherein;

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF_3 , C_1 - C_6 alkoxy, C_2 - C_6 dialkoxymethyl, C_1 - C_6 alkyl, cyano, C_3 - C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkylthio, allyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, aralkoxy, C_2 - C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl, C_1 - C_6 alkylthio, C_1 - C_6 alkylsulfonyl, C_1 - C_6 haloalkylsulfonyl, C_1 - C_6 alkylsulfinyl, C_1 - C_6 haloalkylsulfinyl, arylthio, C_1 - C_6 haloalkoxy, amino, C_1 - C_6 alkylamino, C_2 - C_{15} dialkylamino, hydroxy, carbamoyl, C_1 - C_6 N-alkylcarbamoyl, C_2 - C_{15} N,N-dialkylcarbamoyl, nitro and C_2 - C_{15} dialkylsulfamoyl;

X is $-(\text{CH}_2)_q-$, wherein q is 1-6,

$-(\text{CH}_2)_r-\text{C}\equiv\text{C}-(\text{CH}_2)_r-$,

$-(\text{CH}_2)_r-\text{CH}=\text{CH}-(\text{CH}_2)_r-$,

$-(\text{CH}_2)_r-\overset{\text{O}}{\parallel}{\text{C}}-(\text{CH}_2)_r-$,

$-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-$, wherein each r is independently 0-3 and

wherein Y is O or S; or

C_1 - C_6 alkyl;


Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1 - C_6 alkoxy, C_2 - C_6 dialkoxymethyl, C_1 - C_6 alkyl, cyano, C_3 - C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1 -

-164-

C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

wherein said compound exhibits high binding activity with respect to the sigma receptor.

33. A method of treating a human being suffering from drug abuse, gastrointestinal disorder, or depression, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



wherein n is 0-5;

Cy is C₃-C₈ cycloalkyl;

R is hydrogen or C₁-C₆ alkyl;

R¹ is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O;

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein r is 0-3;

-(CH₂)_r-CH=CH-(CH₂)_r-;

-(CH₂)_r-C(=O)-(CH₂)_r-;

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and

wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen); and

Z is hydrogen, cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆


-165-

haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

wherein said compound exhibits high binding activity with respect to the sigma receptor.

34. The method of claim 33, wherein said compound is selected from the group consisting of N-(3-cyclohexylpropyl)-3-phenylpropylamine, N-cyclohexylmethyl-3-phenylpropylamine, N-(5-cyclohexylpentyl)benzylamine, 5-cyclohexylpentylamine, N-methyl-5-cyclohexylpentylamine, N,N-Dimethyl-5-cyclohexylpentylamine, N-cyclohexylmethyl-5-cyclohexyl-n-pentylamine, and N-cyclohexylmethyl-N-methyl-5-cyclohexyl-n-pentylamine.

35. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



wherein n is 0-5;

Cy is C₃-C₈ cycloalkyl;

R is hydrogen or C₁-C₆ alkyl;


R¹ is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O;

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein r is 0-3;

-(CH₂)_r-CH=CH-(CH₂)_r-;

-166-




$-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S ; or

C_1-C_6 alkyl (wherein Z is hydrogen); and

5 Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted
10 aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkylamino, C_2-C_{15} dialkylamino, hydroxy, carbamoyl, C_1-C_6 N -alkylcarbamoyl, C_2-C_{15} N,N -dialkylcarbamoyl, nitro, C_2-C_{15} dialkylsulfamoyl or methylenedioxy;
15


wherein said compound exhibits high binding activity with respect to the sigma receptor.

20 36. A method of treating a human being suffering from drug abuse, gastrointestinal disorder or depression, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



wherein X^1 is $-(\text{CH}_2)_r-\text{C}\equiv\text{C}-(\text{CH}_2)_r-$, wherein each r is 0-3 independently;

30 $-(\text{CH}_2)_r-\text{CH}=\text{CH}-(\text{CH}_2)_r-$;



$-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-$, wherein Y is O or S ; or

35 C_1-C_6 alkyl (wherein Z is hydrogen);

R^2 is independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6

-167-

dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

V is N or CM, wherein M is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, trifluoromethyl, hydroxy or one half of a double bond with the neighboring endocyclic carbon;

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein r is 0-3;

-(CH₂)_r-CH=CH-(CH₂)_r-;

$$\begin{array}{c} \text{O} \\ \parallel \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-;} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and wherein Y is O or S; or


C₁-C₆ alkyl (wherein Z is hydrogen); and

Z is hydrogen, cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

wherein said compound exhibits high binding activity with respect to the sigma receptor.

-168-

37. A method of treating a human being suffering from a central nervous system disorder, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



10 wherein X^1 is $-(CH_2)_r-C\equiv C-(CH_2)_r-$, wherein each r is 0-3 independently;

$-(CH_2)_r-CH=CH-(CH_2)_r-$;

15 $-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$;
 $-(CH_2)_r-Y-(CH_2)_r-$, wherein Y is O or S; or
 C_1-C_6 alkyl (wherein Z is hydrogen);


R^2 is independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro and C_2-C_{15} dialkylsulfamoyl;

30 V is N or CM, wherein M is hydrogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, fluoro, chloro, bromo, trifluoromethyl, hydroxy or one half of a double bond with the neighboring endocyclic carbon;

X is $-(CH_2)_q-$, wherein q is 1-6;

35 $-(CH_2)_r-C\equiv C-(CH_2)_r-$, wherein r is 0-3;
 $-(CH_2)_r-CH=CH-(CH_2)_r-$;

-169-



$-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S ; or

$\text{C}_1\text{-C}_6$ alkyl (wherein Z is hydrogen); and

5 Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_2\text{-C}_6$ dialkoxymethyl, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_3\text{-C}_{15}$ dialkylaminoalkyl, carboxy, carboxamido, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ haloalkylthio, allyl, aralkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, aroyl, aralkoxy, $\text{C}_2\text{-C}_6$ carboxylic acyl, aryl, substituted
10 aryl, heteroaryl, substituted heteroaryl, $\text{C}_3\text{-C}_6$ heterocycloalkyl, $\text{C}_1\text{-C}_6$ alkylthio, $\text{C}_1\text{-C}_6$ alkylsulfonyl, $\text{C}_1\text{-C}_6$ haloalkylsulfonyl, $\text{C}_1\text{-C}_6$ alkylsulfinyl, $\text{C}_1\text{-C}_6$ haloalkylsulfinyl, arylthio, $\text{C}_1\text{-C}_6$ haloalkoxy, amino, $\text{C}_1\text{-C}_6$ alkylamino, $\text{C}_2\text{-C}_{15}$ dialkylamino, hydroxy, carbamoyl, $\text{C}_1\text{-C}_6$ N -alkylcarbamoyl, $\text{C}_2\text{-C}_{15}$ N,N -dialkylcarbamoyl, nitro and $\text{C}_2\text{-C}_{15}$ dialkylsulfamoyl;

15 wherein said compound exhibits high binding activity with respect to the sigma receptor.

38. A method of treating a human being suffering from a
20 central nervous system disorder, drug abuse, gastrointestinal disorder or depression, which comprises administering to said human a therapeutically effective amount of a compound having the formula:



25 wherein R^5 and R^6 are independently a C_{1-8} alkyl group;


30 R^7 is hydrogen or a C_{1-8} alkyl group substituted by an arylacetoxo or arylcarboxy group; and

X is $-(\text{CH}_2)_q-$, wherein q is 1-6;

$-(\text{CH}_2)_r-\text{C}\equiv\text{C}-(\text{CH}_2)_r-$, wherein r is 0-3;

$-(\text{CH}_2)_r-\text{CH}=\text{CH}-(\text{CH}_2)_r-$;

35



-170-

$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S; or

C_1-C_6 alkyl (wherein R^7 is hydrogen);


wherein said compound exhibits a high binding activity with respect to the sigma receptors.

39. The method of claim 38, wherein said compound is selected from the group consisting of N,N-dimethyl-n-hexylamine, N-methyl-N-propylhexylamine, N,N'-diethyl-2-(diphenylacetoxy)ethylamine, and N,N'-diethyl-2-(9-fluorene-carboxy)ethylamine.

40. A method of treating or preventing a central nervous system disorder, angina, migraine or hypertension, comprising administering to an animal an effective amount of a compound selected from the group consisting of N-phenyl-N'-(3-(1-phthalimido)propyl)piperazine, N-(4-(1-phthalimido)butyl)-N'-(o-methoxyphenyl)piperazine, and N-phenyl-N'-(4-(1-phthalimido)butyl)piperazine.

41. The method of any one of claims 1-3, 5-10, 15, 18, 20, 23, 25, 27, 29, 31, 32, 33, 35-38 and 40 wherein said compound is administered as part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

42. A compound of the formula



wherein:

-171-

R is hydrogen or methyl;

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O; or R and R¹ together form a morpholino ring;

n is 0-5;

W is -(CH₂)_p- or -H H-, wherein p is 1-3;

X is -(CH₂)_q-, wherein q is 3-6;

-(CH₂)_r-C≡C-(CH₂)_r-;

-(CH₂)_r-CH=CH-(CH₂)_r-;

$$\begin{array}{c} \text{O} \\ || \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-;} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and wherein Y is O or S;

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

-172-


with the proviso that when W is -H, H-, R and R¹ are hydrogen and n is 0-2, then q is 4-6;

wherein said compound exhibits high binding activity with respect to the sigma receptor.

5

43. A compound of the formula:

10



wherein:

15

R is hydrogen or C₁-C₆ alkyl;

20

T is cycloalkyl, aryl, heteroaryl, substituted aryl, or substituted heteroaryl, wherein said substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

25

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O;

30

n is 0-5;

X is -(CH₂)_q-, wherein q is 3-6;

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-;

35

$\text{-(CH}_2\text{)}_r\text{-}\overset{\text{O}}{\parallel}\text{C-(CH}_2\text{)}_r\text{-; or}$

$\text{-(CH}_2\text{)}_r\text{-Y-(CH}_2\text{)}_r\text{-, wherein each r is independently 0-3 and}$

wherein Y is O or S;

-173-

Z is cycloalkyl, an arylcarboxy, or an arylacetoxo group, wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylene dioxy;

with the proviso that when R and R¹ are hydrogen and n is 0-2, then q is 4-6;


wherein said compound exhibits high binding activity with respect to the sigma receptor.

44. A compound selected from the group consisting of N-(3-phenylpropyl)-1-(4-propyl-phenyl)-isopropylamine, N-(3-phenylpropyl)-1-(4-benzoxyphe-
nyl)-isopropylamine, N-methyl-N-(3-phenylpropyl)-1-(4-propylphenyl)-isopropylamine, N-(3-phenylpropyl)-1-phenyl-2-pentylamine, N-(4-phenylbutyl)-1-phenyl-2-pentylamine, N-(3-phenylpropyl)-N-(6-phenylhexyl)amine, N-(3-phenylpropyl)-N-(5-phenylpentyl)amine, N-propyl-N-methyl-5-phenylpentylamine, N-methyl-N-(3-phenylpropyl)-1-phenylisopropylamine, N-methyl-N-(3-methyl-2-butenyl)-1-phenylisopropylamine, N-methyl-N-(3-methylbutyl)-1-phenylisopropylamine, N-methyl-N-(3-phenylpropyl)-1-phenyl-2-pentylamine, N-methyl-N-(3-methylbutyl)-1-isopropylamine, N-methyl-N-(3-phenylbutyl)-1-phenyl-2-pentylamine, N-propyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine, N-benzyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine, N-phenyl-(5-phenyl)pentylamine, N-methyl-N-(3-phenylpropyl)-5-phenylpentylamine, N-(2-(o-methylphenyl)ethyl)-5-phenylpentylamine, N-(2-(m-methylphenyl)ethyl)-5-phenylpentylamine, N-(2-(p-methylphenyl)ethyl)-5-phenylpentylamine, N-benzyl-5-phenylpentylamine, N-benzyl-N-methyl-5-

-174-

phenylpentylamine, N-(2-(3-hydroxyphenyl)ethyl)-5-phenylpentylamine, N-(2-(2-hydroxyphenyl)ethyl)-5-phenylpentylamine, N,N'-diethyl-2-(diphenylacetoxy)ethylamine, N,N'-diethyl-2-(9-fluorene-9-carboxy)ethylamine, N,N-Dimethyl-5-phenylpentylamine, N-Benzyl-N-(3-phenylpropyl)-1-phenyl-2-propylamine, N-Benzyl-N-methyl-5-phenylpentylamine, N-Benzyl-5-phenylpentylamine, and N-(2-phenethyl)-N-methylpentylamine.

45. A compound of the formula:



wherein:

R is hydrogen or methyl;

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

R¹ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O; or

R and R¹ together form a morpholino ring;

n is 1-3;

-175-

p is 1-3;

X is $-(CH_2)_q-$, wherein q is 1-6,

$-(CH_2)_r-C\equiv C-(CH_2)_r-$,

$-(CH_2)_r-CH=CH-(CH_2)_r-$,

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$, or


$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-1 and

wherein Y is O or S;

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkylamino, C_2-C_{15} dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro and C_2-C_{15} dialkylsulfamoyl;

wherein said compound exhibits high binding activity with respect to the sigma receptor.

46. A compound of the formula:



wherein:

R^2 is selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxy-methyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6

-176-

heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

X is -(CH₂)_q-, wherein q is 1-6,

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-,

10

$$\begin{array}{c} \text{O} \\ || \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-,} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and

wherein Y is O or S; or

15

C₁-C₆ alkyl (wherein Z is hydrogen);

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy;

20

25


V is N or CM, wherein M is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, hydroxy, fluoro, chloro, bromo or represents one half of a double bond with a neighboring endocyclic carbon;

30

wherein said compound exhibits a high binding activity with respect to the sigma receptor.


-177-

47. The compound claim 46, wherein V is N, said compound having the formula




48. A compound selected from the group consisting of N-(3-trifluoromethylphenyl)-N'-(4-phenylbutyl)piperazine, N-(3-chlorophenyl)-N'-benzylpiperazine, N-(3-chlorophenyl)-N'-(3-phenylpropyl)piperazine, N-phenyl-N'-(3-phenylpropyl)piperazine, N-phenyl-N'-(3-phenylbutyl)piperazine, N-(2-naphthyl)-N'-(3-phenylpropyl)-piperazine, N-phenyl-N'-(3-(2-naphthyl)propyl)-piperazine, N-phenyl-N'-propylpiperazine, N-(4-chlorophenyl)-N'-(3-phenylpropyl)piperazine, N-benzyl-N'-(4-phenylbutyl)-piperazine, N-phenyl-N'-(4-phthalimidobutyl)piperazine, N-phenyl-N'-(5-phthalimidopentyl)piperazine, N-(5-phenylpentyl)-4-benzylpiperidine and N-(5-phenylpentyl)-4-benzyl-4-hydroxy-piperidine.

49. The compound of claim 46, wherein V is CM, said compound having the formula



50. A compound of the formula:



-178-

wherein:

R^3 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_2 - C_6 dialkoxymethyl, C_3 - C_{15} dialkylaminoalkyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, C_2 - C_6 acyl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl;

X is $-(CH_2)_q-$, wherein q is 1-6,

$-(CH_2)_r-C\equiv C-(CH_2)_r-$,

$-(CH_2)_r-CH=CH-(CH_2)_r-$,

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$,

$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and wherein Y is O or S; or

C_1 - C_6 alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1 - C_6 alkoxy, C_2 - C_6 dialkoxymethyl, C_1 - C_6 alkyl, cyano, C_3 - C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkylthio, allyl, aralkyl, C_3 - C_6 cycloalkyl, aroyl, aralkoxy, C_2 - C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3 - C_6 heterocycloalkyl, C_1 - C_6 alkylthio, C_1 - C_6 alkylsulfonyl, C_1 - C_6 haloalkylsulfonyl, C_1 - C_6 alkylsulfinyl, C_1 - C_6 haloalkylsulfinyl, arylthio, C_1 - C_6 haloalkoxy, amino, C_1 - C_6 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C_1 - C_6 N-alkylcarbamoyl, C_2 - C_{15} N,N-dialkylcarbamoyl, nitro, C_2 - C_{15} dialkylsulfamoyl or methylenedioxy;

V is N or CM, wherein M is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, hydroxy, fluoro, chloro, bromo or represents one half of a double bond with a neighboring endocyclic carbon;

wherein said compound exhibits a high binding activity with respect to the sigma receptor.


51. The compound of claim 50 selected from the group consisting of N-methyl-N'-(4-phenyl-3-(E)butenyl)piperazine, N-methyl-N'-(4-phenyl-3-(Z)butenyl)-piperazine, N-methyl-N'-(4-(3-trifluoromethylphenyl)-3-(Z)butenyl)piperazine, N-methyl-N'-(4-phenylbutyl)piperazine, N-benzyl-N'-(4-phthalimidobutyl)-

-179-

piperazine, N-(2-methoxyphenyl)-N'-(4-phthalimidobutyl)-piperazine, N-(5-phenylpentyl)-4-benzylpiperidine, and N-(5-phenylpentyl)-4-benzyl-4-hydroxy-piperidine.

52. A compound selected from the group consisting of N-(3-trifluoromethylphenyl)-N'-(4-phenylbutyl)piperazine, N-(3-chlorophenyl)-N'-(3-phenylpropyl)piperazine, N-phenyl-N'-(3-phenylpropyl)piperazine, N-phenyl-N'-(3-phenylbutyl)piperazine, N-naphthyl-N'-(3-phenylpropyl)-piperazine, N-phenyl-N'-(3-naphthylpropyl)piperazine, N-phenyl-N'-propylpiperazine, N-(4-chlorophenyl)-N'-(3-phenylpropyl)piperazine, N-benzyl-N'-(4-phenylbutyl)piperazine, N-phenyl-N'-(4-phthalimidobutyl)-piperazine, N-phenyl-N'-(5-phthalimidopentyl)piperazine, N-(5-phenylpentyl)-4-benzylpiperidine and N-(5-phenylpentyl)-4-benzyl-4-hydroxy-piperidine..

53. A compound having the formula





wherein

R^2 is selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxy-methyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkylamino, C_2-C_{15} dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro and C_2-C_{15} dialkylsulfamoyl;

X is $-(CH_2)_r-C\equiv C-(CH_2)_r-$,

-180-




5 $-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and Y is O or S;

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro, C_2-C_{15} dialkylsulfamoyl or methylenedioxy;

wherein said compound exhibits a high binding activity with respect to the sigma receptor.

54. A compound of the formula:





30 wherein:


R^4 is hydrogen or an aryl group substituted with a group selected from the group consisting of C_1-C_6 alkyl, C_1-C_6 alkenyl, C_2-C_6 dialkoxymethyl, C_3-C_{15} dialkylaminoalkyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, C_2-C_6 acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl;

R^5 is hydrogen or hydroxy;

X is $-(CH_2)_q-$, wherein q is 1-6,



-181-



-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and
 5 wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted
 by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆
 dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl,
 10 carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio,
 allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆
 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted
 heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkyl-
 sulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆
 15 haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆
 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-
 alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅
 dialkylsulfamoyl or methylenedioxy;

wherein said compound exhibits a high binding activity with
 20 respect to the sigma receptors.


55. The compound of claim 54 selected from the group
 consisting of N-(5-phenylpentyl)piperidine, N-(8-phenylheptyl)-
 piperidine, N-(5-(4-methoxyphenyl)pentyl)piperidine, N-(3-
 25 phenylpropyl)piperidine, N-(5-cyclohexyl)pentylpiperidine
 N-benzylpiperidine, N-(2-phenethyl)-4-hydroxy-4-phenylpiperidine
 N-(2-phenethyl)-4-hydroxy-4-t-butylpiperidine, N-(5-(4-
 chlorophenyl)-5-pentanone-1-yl)piperidine, N-(5-(4-chlorophenyl)-
 5-pentanone-1-yl)-4-phenylpiperidine, N-(5-(4-methoxyphenyl)-5-
 30 pentanone-1-yl)piperidine, N-(5-(4-methoxyphenyl)-5-pentanone-1-
 yl)-4-phenylpiperidine, N-(5-(4-methoxyphenyl)pentyl)-4-
 phenylpiperidine, N-(5-phenyl-5-pentanone-1-yl)-4-phenylpiperidine
 N-(5-(4-chlorophenyl)pentyl)-4-phenylpiperidine, N-(5-(3-
 methoxyphenyl)-5-pentanone-1-yl)piperidine, N-(5-(3-chlorophenyl)-
 5-pentanone-1-yl)piperidine, N-(5-(3-chlorophenyl)-5-pentanone-1-
 35 yl)-4-phenylpiperidine, N-(5-(3-methoxyphenyl)-5-pentanone-1-yl)-

-182-

4-phenylpiperidine, N-(4-(4-fluorophenyl)-4-butanon-1-yl)piperidine, N-(5-(4-fluorophenyl)-5-pentanon-1-yl)piperidine
 N-(5-(4-fluorophenyl)-5-pentanon-1-yl)-4-phenylpiperidine, N-(5-
 5 (4-fluorophenyl)-5-pentanon-1-yl)-4-(3-chlorophenyl)-4-
 hydroxypiperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-
 fluorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-5-
 pentanon-1-yl)-4-(4-fluorophenyl)piperidine, N-(5-(4-chloro-
 phenyl)-5-pentanon-1-yl)-4-(4-fluorophenyl)-1,2,3,6-tetrahydro-
 pyridine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-(4-fluoro-
 10 phenyl)-piperidine, N-(5-(4-chlorophenyl)-5-pentanon-1-yl)-4-
 (chlorophenyl)-1,2,3,6-tetrahydropyridine, N-(5-(4-chlorophenyl)-
 5-pentanon-1-yl)-4-(chlorophenyl)piperidine, N-(5-(3,4-dichloro-
 phenyl)-5-pentanon-1-yl)-4-(chlorophenyl)-piperidine, N-(5-
 cyclopentylpentan-5-on-1-yl)piperidine, N-(5-(3,4-methylene-
 15 dioxyphenyl)penta-2,4-dienyl)piperidine, N-phenyl-(5-
 phenyl)pentylamine, N-methyl-N-(3-phenylpropyl)-5-
 phenylpentylamine, N-benzyl-N-methyl-5-phenylpentylamine, N-(2-
 (o-methoxyphenyl)ethyl)-5-phenylpentylamine, N-(2-(m-
 methoxyphenyl)ethyl)-5-phenylpentylamine, N-(2-(p-
 20 methoxyphenyl)ethyl)-5-phenylpentylamine, N-benzyl-5-
 phenylpentylamine, N-(2-(m-hydroxyphenyl)ethyl)-5-
 phenylpentylamine, and N-(2-(o-hydroxyphenyl)ethyl)-5-
 phenylpentylamine.

25 56. A compound having the formula:


30



wherein,

35 X is $-(CH_2)_q-$, wherein q is 1-6,
 $-(CH_2)_r-C\equiv C-(CH_2)_r-$,
 $-(CH_2)_r-CH=CH-(CH_2)_r-$,

-183-



$$-(\text{CH}_2)_r-\text{Y}-(\text{CH}_2)_r-, \text{ wherein each } r \text{ is independently } 0-3 \text{ and}$$

5 wherein Y is O or S; or

$\text{C}_1\text{-C}_6$ alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_2\text{-C}_6$ dialkoxymethyl, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_3\text{-C}_{15}$ dialkylaminoalkyl, carboxy, carboxamido, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ haloalkylthio, allyl, aralkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, aroyl, aralkoxy, $\text{C}_2\text{-C}_6$ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, $\text{C}_3\text{-C}_6$ heterocycloalkyl, $\text{C}_1\text{-C}_6$ alkylthio, $\text{C}_1\text{-C}_6$ alkylsulfonyl, $\text{C}_1\text{-C}_6$ haloalkylsulfonyl, $\text{C}_1\text{-C}_6$ alkylsulfinyl, $\text{C}_1\text{-C}_6$ haloalkylsulfinyl, arylthio, $\text{C}_1\text{-C}_6$ haloalkoxy, amino, $\text{C}_1\text{-C}_6$ alkyl-amino, dialkylamino, hydroxy, carbamoyl, $\text{C}_1\text{-C}_6$ N-alkylcarbamoyl, $\text{C}_2\text{-C}_{15}$ N,N-dialkylcarbamoyl, nitro, $\text{C}_2\text{-C}_{15}$ dialkylsulfamoyl or methylenedioxy;

10


15

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

20

57. N-(5-phenyl)pentyl-3-azabicyclo[3.2.2]nonane, a compound of claim 56.

25 58. A tropane derivative having the formula:



wherein,

35 R^4 is hydrogen or an aryl group substituted with a group selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ dialkoxymethyl, $\text{C}_3\text{-C}_{15}$ dialkylaminoalkyl, aralkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, aroyl, $\text{C}_2\text{-C}_6$ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, $\text{C}_3\text{-C}_6$ heterocycloalkyl;

-184-

R^5 is hydrogen or hydroxy;

X is $-(CH_2)_q-$, wherein q is 1-6,

$-(CH_2)_r-C\equiv C-(CH_2)_r-$,

$-(CH_2)_r-CH=CH-(CH_2)_r-$,

5

$-(CH_2)_r-\overset{\overset{O}{\parallel}}{C}-(CH_2)_r-$,

$-(CH_2)_r-Y-(CH_2)_r-$, wherein each r is independently 0-3 and

10

wherein Y is O or S; or

C_1-C_6 alkyl (wherein Z is hydrogen);

Z is cycloalkyl, aryl or heteroaryl wherein Z may be substituted by chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro, C_2-C_{15} dialkylsulfamoyl or methylenedioxy;

15

20

25


wherein said compound exhibits a high binding activity with respect to the sigma receptors.

59. N-(5-phenyl)pentyl-4-phenyltropan-4-ol, a compound of claim 58.

30

60. A compound having the formula:

35



40

wherein;

-185-

Ar is aryl, heteroaryl, substituted aryl or substituted heteroaryl, wherein the substituent is selected from the group consisting of chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl;

X is -(CH₂)_q-, wherein q is 1-6,

-(CH₂)_r-C≡C-(CH₂)_r-,

-(CH₂)_r-CH=CH-(CH₂)_r-,

$$\begin{array}{c} \text{O} \\ \parallel \\ \text{-(CH}_2\text{)}_r\text{-C-(CH}_2\text{)}_r\text{-,} \end{array}$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and

wherein Y is O or S; or

C₁-C₆ alkyl;

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro or C₂-C₁₅ dialkylsulfamoyl;

wherein said compound exhibits high binding activity with respect to the sigma receptor.

-186-

61. A compound of the formula



wherein n is 0-5;

10 Cy is C₃-C₈ cycloalkyl; and

R¹ is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, fluoro, chloro, bromo, iodo and =O;

R is hydrogen or C₁-C₆ alkyl;

15 X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein r is 0-3;

-(CH₂)_r-CH=CH-(CH₂)_r-;

20

$$\text{-(CH}_2\text{)}_r\text{-}\overset{\text{O}}{\parallel}\text{C-(CH}_2\text{)}_r\text{-};$$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and

wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen); and

25 Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro, C₂-C₁₅ dialkylsulfamoyl or methylenedioxy.


35

62. A compound selected from the group consisting of N-cyclohexylmethyl-3-phenylpropylamine, N-(5-cyclohexylpentyl)benzylamine, 5-cyclohexylpentylamine, N-methyl-5-cyclohexylpentylamine, N,N-Dimethyl-5-cyclohexylpentylamine, N-

-187-

cyclohexylmethyl-5-cyclohexyl-n-pentylamine, and N-cyclohexylmethyl-N-methyl-5-cyclohexyl-n-pentylamine.

63. A compound of the formula



wherein X^1 is $-(CH_2)_r-C\equiv C-(CH_2)_r-$, wherein each r is 0-3 independently;

$-(CH_2)_r-CH=CH-(CH_2)_r-$;

$-(CH_2)_r-C(=O)-(CH_2)_r-$;

$-(CH_2)_r-Y-(CH_2)_r-$, wherein Y is O or S; or

C_1-C_6 alkyl (wherein Z is hydrogen);

R^2 is independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, iodo, CF_3 , C_1-C_6 alkoxy, C_2-C_6 dialkoxymethyl, C_1-C_6 alkyl, cyano, C_3-C_{15} dialkylaminoalkyl, carboxy, carboxamido, C_1-C_6 haloalkyl, C_1-C_6 haloalkylthio, allyl, aralkyl, C_3-C_6 cycloalkyl, aroyl, aralkoxy, C_2-C_6 carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C_3-C_6 heterocycloalkyl, C_1-C_6 alkylthio, C_1-C_6 alkylsulfonyl, C_1-C_6 haloalkylsulfonyl, C_1-C_6 alkylsulfinyl, C_1-C_6 haloalkylsulfinyl, arylthio, C_1-C_6 haloalkoxy, amino, C_1-C_6 alkyl-amino, dialkylamino, hydroxy, carbamoyl, C_1-C_6 N-alkylcarbamoyl, C_2-C_{15} N,N-dialkylcarbamoyl, nitro and C_2-C_{15} dialkylsulfamoyl; V is N or CM, wherein M is hydrogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, fluoro, chloro, bromo, trifluoromethyl, hydroxy or one half of a double bond with the neighboring endocyclic carbon;

X is $-(CH_2)_q-$, wherein q is 1-6;

$-(CH_2)_r-C\equiv C-(CH_2)_r-$, wherein r is 0-3;

$-(CH_2)_r-CH=CH-(CH_2)_r-$;

$-(CH_2)_r-C(=O)-(CH_2)_r-$;


-188-

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and wherein Y is O or S; or

C₁-C₆ alkyl (wherein Z is hydrogen); and

Z is cycloalkyl which may be substituted by chloro, fluoro, bromo, iodo, CF₃, C₁-C₆ alkoxy, C₂-C₆ dialkoxymethyl, C₁-C₆ alkyl, cyano, C₃-C₁₅ dialkylaminoalkyl, carboxy, carboxamido, C₁-C₆ haloalkyl, C₁-C₆ haloalkylthio, allyl, aralkyl, C₃-C₆ cycloalkyl, aroyl, aralkoxy, C₂-C₆ carboxylic acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, C₃-C₆ heterocycloalkyl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, arylthio, C₁-C₆ haloalkoxy, amino, C₁-C₆ alkylamino, C₂-C₁₅ dialkylamino, hydroxy, carbamoyl, C₁-C₆ N-alkylcarbamoyl, C₂-C₁₅ N,N-dialkylcarbamoyl, nitro and C₂-C₁₅ dialkylsulfamoyl.

64. A compound having the formula:



wherein R⁵ and R⁶ are independently a C₁₋₈ alkyl group; R⁷ is hydrogen or a C₁₋₈ alkyl group substituted by an arylacetoxo or arylcarboxy group; and

X is -(CH₂)_q-, wherein q is 1-6;

-(CH₂)_r-C≡C-(CH₂)_r-, wherein r is 0-3;

-(CH₂)_r-CH=CH-(CH₂)_r-;

$\begin{array}{c} O \\ || \\ -(CH_2)_r-C-(CH_2)_r- \end{array}$

-(CH₂)_r-Y-(CH₂)_r-, wherein each r is independently 0-3 and wherein Y is O or S; or

C₁-C₆ alkyl (wherein R⁷ is hydrogen);

wherein said compound exhibits a high binding activity with respect to the sigma receptors.

65. The compound claim 64 selected from the group consisting of N,N-dimethyl-n-hexylamine, N-methyl-N-

-189-

propylhexylamine, N,N'-diethyl-2-(diphenylacetoxy)ethylamine, and N,N'-diethyl-2-(9-fluorene-carboxy)ethylamine.

5 66. A compound selected from the group consisting of N-phenyl-N'-(3-(1-phthalimido)propyl)piperazine, N-(4-(1-phthalimido)butyl)-N'-(o-methoxyphenyl)piperazine, and N-phenyl-N'-(4-(1-phthalimido)butyl)piperazine.

10 67. A pharmaceutical composition comprising a therapeutically effective amount of the compound of any one of claims 42-46, 48, 52-54, 56, 58 and 60-66 and a pharmaceutically acceptable carrier.